

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
30 March 2006 (30.03.2006)

PCT

(10) International Publication Number  
**WO 2006/033620 A1**

## (51) International Patent Classification:

C07D 401/04 (2006.01) A61P 29/00 (2006.01)  
A61K 31/4184 (2006.01) C07D 235/06 (2006.01)  
A61K 31/4439 (2006.01) C07D 405/12 (2006.01)

(74) Agent: ASTRAZENECA; Global Intellectual Property,  
S-151 85 Södertälje (SE).

## (21) International Application Number:

PCT/SE2005/001364

## (22) International Filing Date:

19 September 2005 (19.09.2005)

## (25) Filing Language:

English

## (26) Publication Language:

English

## (30) Priority Data:

0402284-4 21 September 2004 (21.09.2004) SE

(71) Applicant (for all designated States except US): AS-  
TRAZENECA AB [SE/SE]; S-151 85 Södertälje (SE).

## (72) Inventors; and

(75) Inventors/Applicants (for US only): BESIDSKI, Yev-  
genii [SE/SE]; AstraZeneca R & D Södertälje, S-151 85  
Södertälje (SE). KERS, Inger [SE/SE]; AstraZeneca  
R & D Södertälje, S-151 85 Södertälje (SE). NYLÖF,  
Martin [SE/SE]; AstraZeneca R & D Södertälje, S-151  
85 Södertälje (SE). SLAITAS, Andis [LV/LV]; Syntagon  
Baltic, Vecpabrigs Pinki Babiņas pag., LV-2107 Rīga (LV).

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,  
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,  
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY,  
MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO,  
NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK,  
SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,  
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,  
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,  
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,  
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

## Published:

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

For two-letter codes and other abbreviations, refer to the "Guide-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: NEW HETEROCYCLIC AMIDES



(I)

(57) Abstract: The present invention relates to new  
compounds (I) or salts, solvates or solvated salts thereof,  
processes for their preparation and to new intermediates  
used in the preparation thereof, pharmaceutical compositions  
containing said compounds and to the use of said compounds  
in therapy.

## NEW HETEROCYCLIC AMIDES

## FIELD OF THE INVENTION

The present invention relates to new compounds, to pharmaceutical compositions containing said compounds and to the use of said compounds in therapy. The present invention further relates to processes for the preparation of said compounds and to new intermediates used in the preparation thereof.

## BACKGROUND OF THE INVENTION

Pain sensation in mammals is due to the activation of the peripheral terminals of a specialized population of sensory neurons known as nociceptors. Capsaicin, the active ingredient in hot peppers, produces sustained activation of nociceptors and also produces a dose-dependent pain sensation in humans. Cloning of the vanilloid receptor 1 (VR1 or TRPV1) demonstrated that VR1 is the molecular target for capsaicin and its analogues. (Caterina, M.J., Schumacher, M.A., et al. *Nature* (1997) v.389 p 816-824). Functional studies using VR1 indicate that it is also activated by noxious heat, tissue acidification and other inflammatory mediators (Tominaga, M., Caterina, M.J. et al. *Neuron* (1998) v.21, p.531-543). Expression of VR1 is also regulated after peripheral nerve damage of the type that leads to neuropathic pain. These properties of VR1 make it a highly relevant target for pain and for diseases involving inflammation. While agonists of the VR1 receptor can act as analgesics through nociceptor destruction, the use of agonists, such as capsaicin and its analogues, is limited due to their pungency, neurotoxicity and induction of hypothermia. Instead, agents that block the activity of VR1 should prove more useful. Antagonists would maintain the analgesic properties, but avoid pungency and neurotoxicity side effects.

Compounds with VR1 inhibitor activity are believed to be of potential use for the treatment and/or prophylaxis of disorders such as pain, especially that of inflammatory or traumatic origin such as arthritis, ischaemia, cancer, fibromyalgia, low back pain and post-operative pain (Walker et al *J Pharmacol Exp Ther.* (2003) Jan;304(1):56-62). In addition to this visceral pains such as chronic pelvic pain, cystitis, irritable bowel syndrome (IBS), pancreatitis and the like, as well as neuropathic pain such as sciatica, diabetic neuropathy, HIV neu-

ropathy, multiple sclerosis, and the like (Walker et al *ibid*, Rashid et al J Pharmacol Exp Ther. (2003) Mar;304(3):940-8), are potential pain states that could be treated with VR1 inhibitor. These compounds are also believed to be potentially useful for inflammatory disorders like asthma, cough, inflammatory bowel disease (IBD) (Hwang and Oh Curr Opin Pharmacol (2002) Jun;2(3):235-42). Compounds with VR1 blocker activity are also useful for itch and skin diseases like psoriasis and for gastro-esophageal reflux disease (GERD), emesis, cancer, urinary incontinence and hyperactive bladder (Yiangou et al BJU Int (2001) Jun;87(9):774-9, Szallasi Am J Clin Pathol (2002) 118: 110-21). VR1 inhibitors are also of potential use for the treatment and/or prophylaxis of the effects of exposure to VR1 activators like capsaicin or tear gas, acids or heat (Szallasi *ibid*).


A further potential use relates to the treatment of tolerance to VR1 activators.

VR1 inhibitors may also be useful in the treatment of interstitial cystitis and pain related to interstitial cystitis.

## DETAILED DESCRIPTION OF THE INVENTION

The object of the present invention is to provide compounds exhibiting an inhibitory activity at the vanilloid receptor 1 (VR1).

The present invention provides compounds of formula I



(I)

wherein:

R<sup>1</sup> is H, NO<sub>2</sub>, halo, NR<sup>6</sup>R<sup>7</sup>, C<sub>1-6</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>1-6</sub>haloalkyl, C<sub>1-6</sub>haloalkylO, R<sup>6</sup>OC<sub>0-6</sub>alkyl, R<sup>6</sup>CO, R<sup>6</sup>OCO or CONR<sup>6</sup>R<sup>7</sup>;  
m is 0, 1, 2 or 3;

$R^2$  is H,  $NO_2$ , halo,  $NR^6R^7$ ,  $C_{1-6}alkyl$ ,  $C_{2-6}alkenyl$ ,  $C_{2-6}alkynyl$ ,  $C_{1-6}haloalkyl$ ,  $C_{1-6}haloalkylO$ , cyano,  $R^6OC_{0-6}alkyl$ ,  $R^6CO$ ,  $R^6OCO$ ,  $R^6CONR^7$ ,  $R^6R^7NCO$ ,  $R^8SO_2$ ,  $R^8SO_2HN$ ,  $arylC_{0-6}alkyl$  or  $heteroarylC_{0-6}alkyl$ ;

$R^3$  and  $R^5$  are each independently H or  $C_{1-4}alkyl$ ;

5  $R^2$  and  $R^3$  optionally form a ring;

p is 0, 1 or 2;

n is 0, 2, 3 or 4;

$R^5$  is  $C_{1-10}alkyl$ ,  $C_{6-10}arylC_{0-6}alkyl$ ,  $C_{3-7}cycloalkylC_{0-6}alkyl$  or  $C_{5-6}heteroarylC_{0-6}alkyl$ , whereby any aryl, heteroaryl or cycloalkyl may be fused with aryl, heteroaryl,

10  $C_{3-7}cycloalkyl$  or  $C_{3-7}heterocycloalkyl$ , and which  $R^5$  may be substituted with one or more A;

A is H, OH,  $NO_2$ , cyano,  $R^6CO$ ,  $R^6O(CO)$ , halo,  $C_{1-6}alkyl$ ,  $NR^6R^7$ ,  $C_{1-6}haloalkyl$ ,  $C_{1-6}haloalkylO$ ,  $R^6OC_{0-6}alkyl$ , hydroxy $C_{1-6}alkyl$ ,  $R^8SO_2$ ,  $R^8SO_2HN$ ,  $C_{5-6}arylO$  or  $CONR^6R^7$ ;

15  $R^6$  and  $R^7$  are each independently H or  $C_{1-6}alkyl$ ; and

$R^8$  is  $NR^6R^7$  or  $C_{1-4}alkyl$

or salts, solvates or solvated salts thereof.

20 Listed below are definitions of various terms used in the specification and claims to describe the present invention.

For the avoidance of doubt it is to be understood that where in this specification a group is qualified by 'hereinbefore defined', 'defined hereinbefore' or 'defined above' the said group encompasses the first occurring and broadest definition as well as each and all of the  
25 other definitions for that group.

For the avoidance of doubt it is to be understood that in this specification 'C<sub>1-6</sub>' means a carbon group having 1, 2, 3, 4, 5 or 6 carbon atoms.

30 In this specification, unless stated otherwise, the term "alkyl" includes both straight and branched chain alkyl groups and may be, but are not limited to methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl, n-pentyl, i-pentyl, t-pentyl, neo-pentyl, n-hexyl or

i-hexyl, t-hexyl. The term  $C_{1-3}$  alkyl having 1 to 3 carbon atoms and may be methyl, ethyl, n-propyl, i-propyl or *tert*-butyl.

The term 'C<sub>0</sub>' means a bond or does not exist. For example when R<sup>3</sup> is C<sub>0</sub>alkyl, R<sup>3</sup> is a bond and "arylC<sub>0</sub>alkyl" is equivalent with "aryl", "C<sub>2</sub>alkylOC<sub>0</sub>alkyl" is equivalent with "C<sub>2</sub>alkylO".

In this specification, unless stated otherwise, the term "alkenyl" includes both straight and branched chain alkenyl groups. The term "C<sub>2-6</sub>alkenyl" having 2 to 6 carbon atoms and one or two double bonds, may be, but is not limited to vinyl, allyl, propenyl, butenyl, crotyl, pentenyl, or hexenyl, and a butenyl group may for example be buten-2-yl, buten-3-yl or buten-4-yl.

In this specification, unless stated otherwise, the term "alkynyl" includes both straight and branched chain alkynyl groups. The term "C<sub>2-6</sub>alkynyl" having 2 to 6 carbon atoms and one or two trippel bonds, may be, but is not limited to etynyl, propargyl, pentynyl or hexynyl and a butynyl group may for example be butyn-3-yl or butyn-4-yl.

In this specification, unless stated otherwise, the term "cycloalkyl" refers to an optionally substituted, saturated cyclic hydrocarbon ring system. The term "C<sub>3-7</sub>cycloalkyl" may be cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl.

The term "heterocycloalkyl" denotes a 3- to 7-membered, non-aromatic, partially or completely saturated hydrocarbon group, which contains one ring and at least one heteroatom. Examples of said heterocycle include, but are not limited to pyrrolidiny, pyrrolidonyl, piperidiny, piperaziny, morpholinyl, oxazolyl, 2-oxazolidonyl or tetrahydrofuranlyl.

In this specification, unless stated otherwise, the term "aryl" refers to an optionally substituted monocyclic or bicyclic hydrocarbon unsaturated aromatic ring system. Examples of "aryl" may be, but are not limited to phenyl and naphthyl.

In this specification, unless stated otherwise, the term "heteroaryl" refers to an optionally substituted monocyclic or bicyclic unsaturated aromatic ring system containing at least one heteroatom selected independently from N, O or S. Examples of "heteroaryl" may be, but are not limited to pyridyl, pyrrolyl, furyl, thienyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, pyrazolyl, benzofuryl, indolyl, isoindolyl, benzimidazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, tetrazolyl, triazolyl or oxazolyl.

In this specification, unless stated otherwise, the terms "arylalkyl" and "heteroarylalkyl" refer to a substituent that is attached via the alkyl group to an aryl or heteroaryl group.

In this specification, unless stated otherwise, the terms "halo" and "halogen" may be fluoro, iodo, chloro or bromo.

In this specification, unless stated otherwise, the term "haloalkyl" means an alkyl group as defined above, which is substituted with halo as defined above. The term "C<sub>1-6</sub>haloalkyl" may include, but is not limited to fluoromethyl, difluoromethyl, trifluoromethyl, fluoroethyl, difluoroethyl or bromopropyl. The term "C<sub>1-6</sub>haloalkylo" may include, but is not limited to fluoromethoxy, difluoromethoxy, trifluoromethoxy, fluoroethoxy or difluoroethoxy.

The present invention provides compounds selected from the group consisting of  
N-[3-[2-(dimethylamino)ethoxy]phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-(methoxymethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-(1,3-dihydro-2-benzofuran-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-methoxy-5-(methoxymethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-(methoxymethyl)-5-(trifluoromethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-[3-(methoxymethyl)-5-(trifluoromethyl)phenyl]acetamide,  
N-[3-cyano-5-(trifluoromethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-acetyl-5-(trifluoromethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-[3-acetyl-5-(trifluoromethyl)phenyl]acetamide,

- 2-(7-acetyl-1H-benzimidazol-1-yl)-N-[3-cyano-5-(trifluoromethyl)phenyl]acetamide,  
N-[3-(1-methoxyethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-chloro-6-methoxy-1H-benzimidazol-1-yl)-N-(2,3-dihydro-1H-inden-5-yl)acetamide,  
N-[3-(2-methoxyethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
5 N-[3-methoxy-5-(2-methoxyethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-(2-methoxyethoxy)-5-(trifluoromethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-methoxy-5-(tetrahydrofuran-2-ylmethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
10 N-[3-methoxy-5-(tetrahydrofuran-3-yloxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-methoxy-5-(trifluoromethyl)phenyl]-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-4-carboxamide,  
2-(7-amino-1H-benzimidazol-1-yl)-N-(4-tert-butylbenzyl)acetamide,  
15 N-(4-tert-butylbenzyl)-2-(7-iodo-1H-benzimidazol-1-yl)acetamide,  
2-[7-(1-hydroxy-1-methylethyl)-1H-benzimidazol-1-yl]-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,  
N-(4-tert-butylbenzyl)-2-[7-(1-hydroxyethyl)-1H-benzimidazol-1-yl]acetamide,  
N-(2,3-dihydro-1H-inden-5-yl)-2-(7-pyridin-2-yl-1H-benzimidazol-1-yl)acetamide,  
20 N-(4-tert-butylbenzyl)-2-(7-pyridin-2-yl-1H-benzimidazol-1-yl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-(4-tert-butylbenzyl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,  
2-(7-nitro-1H-benzimidazol-1-yl)-N-(3,4,5-trifluorophenyl)acetamide,  
N-(4-tert-butylbenzyl)-2-(7-cyano-1H-benzimidazol-1-yl)acetamide,  
25 2-(7-cyano-1H-benzimidazol-1-yl)-N-(3,4,5-trimethoxyphenyl)acetamide,  
N-(4-bromo-2-fluorophenyl)-2-(7-cyano-1H-benzimidazol-1-yl)acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-(3,4-difluorophenyl)acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-(3-ethoxyphenyl)acetamide,  
2-(7-nitro-1H-benzimidazol-1-yl)-N-(3,4,5-trimethoxybenzyl)acetamide,  
30 N-(3,4-difluorobenzyl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[2-(4-methoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[2-(3-fluorophenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,

- N-[2-(3-methoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-nitro-1H-benzimidazol-1-yl)-N-{2-[3-(trifluoromethyl)phenyl]ethyl}acetamide,  
N-[2-(3,4-dimethoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-(3,4,5-trimethoxyphenyl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-(3,4-difluorophenyl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-(3,5-dimethoxyphenyl)acetamide,  
N-[2-(3,5-dimethoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-(2,3-dihydro-1H-inden-2-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[2-(5-bromo-2-methoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[1-(4-chlorobenzyl)-2-hydroxyethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-(2-hydroxy-2-phenylethyl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylic acid,  
1-[2-(2,3-dihydro-1H-inden-5-ylamino)-2-oxoethyl]-1H-benzimidazole-7-carboxylic acid,  
N-(3,5-dimethoxyphenyl)-2-[7-(hydroxymethyl)-1H-benzimidazol-1-yl]acetamide,  
1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-N-ethyl-1H-benzimidazole-7-carbox-  
amide,  
1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-N-methyl-1H-benzimidazole-7-carbox-  
amide,  
1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-N,N-dimethyl-1H-benzimidazole-7-car-  
boxamide,  
1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-N-methoxy-1H-benzimidazole-7-car-  
boxamide,  
ethyl 1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylate,  
ethyl 1-{2-[(4-tert-butylbenzyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylate,  
ethyl 1-[2-(2,3-dihydro-1H-inden-5-ylamino)-2-oxoethyl]-1H-benzimidazole-7-carboxy-  
late,  
N-(4-tert-butylbenzyl)-2-[7-(dimethylamino)-1H-benzimidazol-1-yl]acetamide,  
N-(4-methoxy-2-naphthyl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(1H-benzimidazol-1-yl)-N-(4-tert-butylbenzyl)acetamide, and  
2-(1H-benzimidazol-1-yl)-N-(2,3-dihydro-1H-inden-5-yl)acetamide,  
or salts, solvates or solvated salts thereof.



- The present invention further provides compounds selected from the group consisting of
- N-(3,5-dimethoxyphenyl)-2-(7-ethynyl-1H-benzimidazol-1-yl)acetamide,
- N-(3,5-dimethoxyphenyl)-2-(7-prop-1-yn-1-yl-1H-benzimidazol-1-yl)acetamide,
- N-(3,5-dimethoxyphenyl)-2-[7-(1H-1,2,3-triazol-4-yl)-1H-benzimidazol-1-yl]acetamide,,
- 5 N-(3,5-dimethoxyphenyl)-2-[7-(1-methyl-1H-1,2,3-triazol-4-yl)-1H-benzimidazol-1-yl]acetamide,
- N-(3,5-dimethoxyphenyl)-2-[7-(1-methyl-1H-tetrazol-5-yl)-1H-benzimidazol-1-yl]acetamide,
- 2-(7-ethyl-1H-benzimidazol-1-yl)-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,
- 10 2-[7-(2-hydroxyethyl)-1H-benzimidazol-1-yl]-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,
- 2-[7-(2-hydroxy-1-methylethyl)-1H-benzimidazol-1-yl]-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,
- N-[3-methoxy-5-(trifluoromethyl)phenyl]-2-(7-vinyl-1H-benzimidazol-1-yl)acetamide,
- 15 2-(7-isopropenyl-1H-benzimidazol-1-yl)-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide ,
- 2-(7-isopropyl-1H-benzimidazol-1-yl)-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,
- N-(3,5-dimethoxyphenyl)-2-(7-methoxy-1H-benzimidazol-1-yl)acetamide,
- N-(3,5-dimethoxyphenyl)-2-(7-ethoxy-1H-benzimidazol-1-yl)acetamide,
- 20 N-(3,5-dimethoxyphenyl)-2-(7-isopropoxy-1H-benzimidazol-1-yl)acetamide,
- 2-(7-tert-butoxy-1H-benzimidazol-1-yl)-N-(3,5-dimethoxyphenyl)acetamide,
- N-(3,5-dimethoxyphenyl)-2-[7-(trifluoromethoxy)-1H-benzimidazol-1-yl]acetamide
- N-(3,5-dimethoxyphenyl)-2-[7-(methylsulfinyl)-1H-benzimidazol-1-yl]acetamide,
- 2-[7-(difluoromethyl)-1H-benzimidazol-1-yl]-N-(3,5-dimethoxyphenyl)acetamide,
- 25 2-[7-(cyanomethyl)-1H-benzimidazol-1-yl]-N-(3,5-dimethoxyphenyl)acetamide,
- 2-[7-(aminomethyl)-1H-benzimidazol-1-yl]-N-(3,5-dimethoxyphenyl)acetamide
- N-(3,5-dimethoxyphenyl)-2-{7-[(dimethylamino)methyl]-1H-benzimidazol-1-yl}acetamide,
- 2-(7-cyclopropyl-1H-benzimidazol-1-yl)-N-(3,5-dimethoxyphenyl)acetamide,
- 30 2-(7-cyclobutyl-1H-benzimidazol-1-yl)-N-(3,5-dimethoxyphenyl)acetamide,
- N-(3,5-dimethoxyphenyl)-2-[7-(methoxymethyl)-1H-benzimidazol-1-yl]acetamide,
- N-(1-isopropyl-1H-benzimidazol-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,

- 2-(7-fluoro-1H-benzimidazol-1-yl)-N-(1-isopropyl-1H-benzimidazol-5-yl)acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-(1-isopropyl-1H-benzimidazol-5-yl)acetamide,  
N-(1-tert-butyl-1H-benzimidazol-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-(1-tert-butyl-2-methyl-1H-benzimidazol-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-(1-isopropyl-2-methyl-1H-benzimidazol-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-(1-isopropyl-2-methyl-1H-benzimidazol-5-yl)acetamide,  
N-(1-tert-butyl-2-methyl-1H-benzimidazol-5-yl)-2-(7-cyano-1H-benzimidazol-1-yl)acetamide,  
N-(1-tert-butyl-1H-benzimidazol-5-yl)-2-(7-cyano-1H-benzimidazol-1-yl)acetamide,  
N-(1-tert-butyl-1H-benzimidazol-5-yl)-2-(7-fluoro-1H-benzimidazol-1-yl)acetamide,  
N-(1-tert-butyl-2-methyl-1H-benzimidazol-5-yl)-2-(7-fluoro-1H-benzimidazol-1-yl)acetamide,  
2-(7-fluoro-1H-benzimidazol-1-yl)-N-(1-isopropyl-2-methyl-1H-benzimidazol-5-yl)acetamide,  
N-[1-isopropyl-7-(trifluoromethyl)-1H-benzimidazol-5-yl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-fluoro-1H-benzimidazol-1-yl)-N-[1-isopropyl-7-(trifluoromethyl)-1H-benzimidazol-5-yl]acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-[1-isopropyl-7-(trifluoromethyl)-1H-benzimidazol-5-yl]acetamide,  
N-2-naphthyl-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-2-naphthylacetamide,  
2-(7-fluoro-1H-benzimidazol-1-yl)-N-2-naphthylacetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-(4-methoxy-2-naphthyl)acetamide,  
2-(7-fluoro-1H-benzimidazol-1-yl)-N-(4-methoxy-2-naphthyl)acetamide,  
N-[3-methoxy-5-(tetrahydro-2H-pyran-2-ylmethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-(2-isopropoxyethoxy)-5-methoxyphenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,

N-[3,5-bis(2-ethoxyethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-{3-methoxy-5-[2-(2-oxopyrrolidin-1-yl)ethoxy]phenyl}-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-methoxy-5-(3-morpholin-4-ylpropoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N,N-diethyl-2-{3-methoxy-5-[[{(7-nitro-1H-benzimidazol-1-yl)acetyl]amino}phenoxy]acetamide,  
N-{3-methoxy-5-[(1-methylpiperidin-2-yl)methoxy]phenyl}-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-{3-[2-(1H-imidazol-1-yl)ethoxy]-5-methoxyphenyl}-2-(7-nitro-1H-benzimidazol-1-yl)acetamide, and  
N-{3-methoxy-5-[(1-methyl-1H-imidazol-5-yl)methoxy]phenyl}-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
or salts, solvates or solvated salts thereof.

The present invention relates to the compounds of the invention as hereinbefore defined as well as to the salts, solvates or solvated salts thereof. Salts for use in pharmaceutical compositions will be pharmaceutically acceptable salts, but other salts may be useful in the production of the compounds of the invention.

A suitable pharmaceutically acceptable salt of the compounds of the invention is, for example, an acid-addition salt, for example an inorganic or organic acid. In addition, a suitable pharmaceutically acceptable salt of the compounds of the invention is an alkali metal salt, an alkaline earth metal salt or a salt with an organic base.

Other pharmaceutically acceptable salts and methods of preparing these salts may be found in, for example, Remington's Pharmaceutical Sciences (18<sup>th</sup> Edition, Mack Publishing Co.).

Some compounds of the invention may have chiral centres and/or geometric isomeric centres (E- and Z- isomers), and it is to be understood that the invention encompasses all such optical, diastereoisomeric and geometric isomers.

The invention also relates to any and all tautomeric forms of the compounds of the invention.

### **Methods of Preparation**

Some compounds of the present invention may be prepared according to the methods described in PCT/SE2004/000738.

Another aspect of the present invention provides processes for preparing compounds of formula I, or salts, solvates or solvated salts thereof.


Throughout the following description of such processes it is to be understood that, where appropriate, suitable protecting groups will be added to, and subsequently removed from, the various reactants and intermediates in a manner that will be readily understood by one skilled in the art of organic synthesis. Conventional procedures for using such protecting groups as well as examples of suitable protecting groups are described, for example, in "Protective Groups in Organic Synthesis", T.W. Green, P.G.M. Wuts, Wiley-Interscience, New York, (1999). References and descriptions of other suitable reactions are described in textbooks of organic chemistry, for example, "Advanced Organic Chemistry", March, 4<sup>th</sup> ed. McGraw Hill (1992) or, "Organic Synthesis", Smith, McGraw Hill, (1994). For representative examples of heterocyclic chemistry see for example "Heterocyclic Chemistry", J. A. Joule, K. Mills, G. F. Smith, 3<sup>rd</sup> ed. Chapman and Hall (1995), p. 189-224 and "Heterocyclic Chemistry", T. L. Gilchrist, 2<sup>nd</sup> ed. Longman Scientific and Technical (1992), p. 248-282.

The term "room temperature" and "ambient temperature" shall mean, unless otherwise specified, a temperature between 16 and 25 °C.

### **Methods of Preparation**

One embodiment of the invention relates to processes for the preparation of the compound of formula I according to Methods A and B, wherein R<sup>1</sup> to R<sup>9</sup>, unless otherwise specified, are defined as in formula I, comprising;

## Method A




whereby the target compound of formula Ia is obtained from the acid of formula II or its deprotonated form, *via* its conversion into an activated form, i.e. either the acyl chloride by treatment with oxalyl chloride or the mixed anhydride by treatment with *O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate and further treatment with an appropriate amine  $\text{NH}_2\text{R}^5$ . This reaction may be performed in any manner known to the skilled man in the art. The activation may be performed using any other similar activating reagent like 1,3-dicyclohexylcarbodiimide, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride or 1,1'-carbonyldiimidazole. Suitable solvents to be used for this reaction may be halogenated hydrocarbons such as chloroform, dichloromethane and dichloroethane or aromatic and heteroaromatic compounds such as benzene, toluene, xylene, pyridine and lutidine or ethers such as ethyl ether, tetrahydrofuran and dioxan or aprotic polar solvents like acetonitrile and dimethylformamide, or any mixtures thereof. Catalysts such as heteroaromatic bases like pyridine and lutidine or tertiary amines like triethylamine, *N*-methylmorpholine and ethyl diisopropylamine may be used as well. The temperature may be between -30 and 50°C and the reaction time between 1 and 30 h. Starting materials, the acids of formula II, may be obtained using multistep procedures described in detail in the following examples of synthesis, starting from commercially available appropriately 1,2,3-trisubstituted benzenes.

Or,

## Method B

whereby the target compound of formula I is obtained from another compound of formula I by a chemical modification of the  $R^2$  substituent using standard methods described in the literature, for example:



wherein, the target compound of formula I is obtained from an amidoalkylbromide and an appropriately substituted benzimidazole. The amidoalkylbromides mentioned may be obtained by amination of the corresponding carboxyalkyl bromides or their acyl chloride derivatives.

Generally, this method yields a mixture of two regio-isomers, which can be separated by use of chromatography. Suitable solvents to be used for this reaction may be tertiary amides such as dimethylformamide or dimethylacetamide or aromatic compounds such as benzene, toluene and xylene, or ethers such as ethyl ether, tetrahydrofuran and dioxan or  
5 alcohols such as methanol, ethanol and propanol, or any mixtures thereof. Bases such as potassium *tert*-butoxide, sodium methoxide and sodium hydride or tertiary amines like triethylamine, *N*-methylmorpholine and ethyl diisopropylamine may be used as well. The temperature may be between 0 and 100°C and the reaction time between 1 and 30 h.

### 10 Intermediates

A further embodiment of the invention relates to compounds selected from the group consisting of

3-methoxy-5-(methoxymethyl)aniline,  
15 3-(methoxymethyl)-5-(trifluoromethyl)aniline,  
1-(methoxymethyl)-3-nitro-5-(trifluoromethyl)benzene,  
1-[3-amino-5-(trifluoromethyl)phenyl]ethanone,  
(7-chloro-6-methoxy-1H-benzimidazol-1-yl)acetic acid,  
2-[(2-chloro-3-methoxy-6-nitrophenyl)amino]ethanol,  
20 2-(7-chloro-6-methoxy-1H-benzimidazol-1-yl)ethanol,  
3-(2-methoxyethoxy)-5-(trifluoromethyl)aniline,  
1-(2-methoxyethoxy)-3-nitro-5-(trifluoromethyl)benzene,  
3-methoxy-5-(tetrahydrofuran-2-ylmethoxy)aniline,  
2-[(3-methoxy-5-nitrophenoxy)methyl]tetrahydrofuran,  
25 3-methoxy-5-(tetrahydrofuran-3-yloxy)aniline,  
3-(3-methoxy-5-nitrophenoxy)tetrahydrofuran,  
5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-4-carboxylic acid,  
methyl 8-amino-1,2,3,4-tetrahydroquinoline-2-carboxylate,  
(7-pyridin-2-yl-1H-benzimidazol-1-yl)acetic acid,  
30 methyl (7-bromo-1H-benzimidazol-1-yl)acetate,  
methyl (7-pyridin-2-yl-1H-benzimidazol-1-yl)acetate,  
3-methoxy-5-(tetrahydro-2H-pyran-2-ylmethoxy)aniline, and

3-(2-isopropoxyethoxy)-5-methoxyaniline

Another embodiment relates to the use of these compounds as intermediates in the preparation of compounds of the invention.

### **Pharmaceutical composition**

According to one embodiment of the present invention there is provided a pharmaceutical composition comprising as active ingredient a therapeutically effective amount of the compound of the invention, or salts, solvates or solvated salts thereof, in association with one or more pharmaceutically acceptable diluents, excipients and/or inert carriers.

The composition may be in a form suitable for oral administration, for example as a tablet, pill, syrup, powder, granule or capsule, for parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion) as a sterile solution, suspension or emulsion, for topical administration e.g. as an ointment, patch or cream or for rectal administration e.g. as a suppository.

In general the above compositions may be prepared in a conventional manner using one or more conventional excipients, pharmaceutical acceptable diluents and/or inert carriers.

Suitable daily doses of the compounds of the invention in the treatment of a mammal, including man, are approximately 0.01 to 250 mg/kg bodyweight at peroral administration and about 0.001 to 250 mg/kg bodyweight at parenteral administration.

The typical daily dose of the active ingredient varies within a wide range and will depend on various factors such as the relevant indication, severity of the illness being treated, the route of administration, the age, weight and sex of the patient and the particular compound being used, and may be determined by a physician.

Examples of pharmaceutical composition



The following illustrate representative pharmaceutical dosage forms containing a compound of the invention, or salts, solvates or solvated salts thereof, (hereafter compound X), for preventive or therapeutic use in mammals:

(a): Tablet	mg/tablet
Compound X	100
Lactose	182.75
Croscarmellose sodium	12.0
Maize starch paste (5% w/v paste)	2.25
Magnesium stearate	3.0

5

(b): Capsule	mg/capsule
Compound X	10
Lactose	488.5
Magnesium stearate	1.5

(c): Injection	(50 mg/ml)
Compound X	5.0% w/v
1M Sodium hydroxide solution	15.0% v/v
0.1M Hydrochloric acid	(to adjust pH to 7.6)
Polyethylene glycol 400	4.5% w/v
Water for injection	up to 100%

The above compositions may be obtained by conventional procedures well known in the pharmaceutical art.

10

### Medical use

Surprisingly, it has been found that the compounds according to the present invention are useful in therapy. The compounds of the invention, or salts, solvates or solvated salts

thereof, as well as their corresponding active metabolites, exhibit a high degree of potency and selectivity for individual vanilloid receptor 1 (VR1) groups. Accordingly, the compounds of the present invention are expected to be useful in the treatment of conditions associated with excitatory activation of vanilloid receptor 1 (VR1).

- 5 The compounds may be used to produce an inhibitory effect of VR1 in mammals, including man.

VR1 are highly expressed in the peripheral nervous system and in other tissues. Thus, it is expected that the compounds of the invention are well suited for the treatment of VR1 mediated disorders.

10

The compounds of the invention are expected to be suitable for the treatment of acute and chronic pain, acute and chronic neuropathic pain and acute and chronic inflammatory pain.

- Examples of such disorder may be selected from the group comprising low back pain, post-operative pain, visceral pains like chronic pelvic pain and the like.

Further relevant disorders may be selected from the group comprising cystitis, including interstitial cystitis and pain related thereto, ischemic, sciatica, diabetic neuropathy, multiple sclerosis, arthritis, fibromyalgia, psoriasis, cancer, emesis, urinary incontinence, hyperactive bladder and HIV neuropathy.

- 20 Additional relevant disorders may be selected from the group comprising gastro-esophageal reflux disease (GERD), irritable bowel syndrome (IBS), inflammatory bowel disease (IBD) and pancreatitis.

Other relevant disorders are related to respiratory diseases and may be selected from the group comprising asthma, cough, chronic obstructive lung disease and emphysema, lung fibrosis and interstitial lung disease.

The VR1 inhibitor(s) may be administered by either an oral or inhaled route. The respiratory disease may be an acute and chronic illness and may be related to infection(s) and/or exposure to environmental pollution and/or irritants.

- The compounds of the invention may also be used as antitoxin to treat (over-) exposure to VR1 activators like capsaicin, tear gas, acids or heat. Regarding heat, there is a potential use for VR1 antagonists in (sun-) burn induced pain, or inflammatory pain resulting from burn injuries.

The compounds may further be used for treatment of tolerance to VR1 activators.

One embodiment of the invention relates to the compounds of the invention as hereinbefore defined, for use as a medicament.

5

Another embodiment of the invention relates to the compounds of the invention as hereinbefore defined, for use as a medicament for treatment of VR1 mediated disorders.

10

A further embodiment of the invention relates to the compounds of the invention as hereinbefore defined, for use as a medicament for treatment of acute and chronic pain disorders.

15

Yet another embodiment of the invention relates to the compounds of the invention as hereinbefore defined, for use as a medicament for treatment of acute and chronic neuropathic pain.

20

Yet a further embodiment of the invention relates to the compounds of the invention as hereinbefore defined, for use as a medicament for treatment of acute and chronic inflammatory pain.

25

One embodiment of the invention relates to the compounds of the invention as hereinbefore defined, for use as a medicament for treatment of low back pain, post-operative pain and visceral pains like chronic pelvic pain.

30

Another embodiment of the invention relates to the compounds of the invention as hereinbefore defined, for use as a medicament for treatment of cystitis, including interstitial cystitis and pain related thereto, ischeamic, sciatica, diabetic neuropathy, multiple sclerosis, arthritis, fibromyalgia, psoriasis, cancer, emesis, urinary incontinence, hyperactive bladder and HIV neuropathy.

A further embodiment of the invention relates to the compounds of the invention as hereinbefore defined, for use as a medicament for treatment of gastro-esophageal reflux disease

(GERD), irritable bowel syndrome (IBS), inflammatory bowel disease (IBD) and pancreatitis.

Yet a further embodiment of the invention relates to the compounds of the invention as  
5 hereinbefore defined, for use as a medicament for treatment of respiratory diseases selected from the group comprising asthma, cough, chronic obstructive lung disease and emphysema, lung fibrosis and interstitial lung disease.

One embodiment of the invention relates to the use of the compound of the invention as  
10 hereinbefore defined, in the manufacture of a medicament for treatment of VR1 mediated disorders and for treatment of acute and chronic pain disorders, acute and chronic neuropathic pain and acute and chronic inflammatory pain, and respiratory diseases and any other disorder mentioned above.

15 Another embodiment of the invention relates to a method of treatment of VR1 mediated disorders and acute and chronic pain disorders, acute and chronic neuropathic pain and acute and chronic inflammatory pain, and respiratory diseases, and any other disorder mentioned above, comprising administering to a mammal, including man in need of such treatment, a therapeutically effective amount of the compounds of the invention, as herein-  
20 before defined.

A further embodiment of the invention relates to a pharmaceutical composition comprising a compound of the invention as hereinbefore defined, for use in treatment of VR1 mediated disorders and for treatment of acute and chronic pain disorders, acute and chronic  
25 neuropathic pain and acute and chronic inflammatory pain, and respiratory diseases, and any other disorder mentioned above.

In the context of the present specification, the term "therapy" and "treatment" includes prevention and prophylaxis, unless there are specific indications to the contrary. The terms  
30 "treat", "therapeutic" and "therapeutically" should be construed accordingly.

In this specification, unless stated otherwise, the term “inhibitor” and “antagonist” mean a compound that by any means, partly or completely, blocks the transduction pathway leading to the production of a response by the ligand.

- 5 The term “disorder”, unless stated otherwise, means any condition and disease associated with vanilloid receptor activity.

### Non- Medical use

- 10 In addition to their use in therapeutic medicine, the compounds of the invention, or salts, solvates or solvated salts thereof, are also useful as pharmacological tools in the development and standardisation of *in vitro* and *in vivo* test systems for the evaluation of the effects of inhibitors of VR1 related activity in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as part of the search for new therapeutics agents.

15

### Examples

The invention will now be illustrated by the following non-limiting examples.

### **Abbreviations**

- 20 DCE dichloroethane  
DCM dichloromethane  
DMAP dimethylaminopyridine  
EDC 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride  
HATU O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium  
25 hexafluorophosphate  
HPLC high performance liquid chromatography  
LC liquid chromatography  
MS mass spectrometry  
ret. time retention time  
30 TFA trifluoroacetic acid  
THF tetrahydrofuran  
DMF dimethylformamide

TMEDA tetramethylethylenediamine

EtOAc ethyl acetate

### General methods

5

All starting materials are commercially available or described in the literature. The  $^1\text{H}$  NMR spectra were recorded on Bruker at 400 MHz. The mass spectra were recorded utilising electrospray (LC-MS; LC: Waters 2790, column XTerra MS  $\text{C}_8$  2.5  $\mu\text{m}$  2.1X30 mm, buffer gradient  $\text{H}_2\text{O}+0.1\%\text{TFA}:\text{CH}_3\text{CN}+0.04\%\text{TFA}$ , MS: micromass ZMD// ammonium acetate buffer) ionisation techniques.

10

### Synthesis of the intermediates: 7- substituted 1*H*-benzimidazol-1-yl-acetic acids, 1) thru 7)

#### 15 1) (7-Nitro-1*H*-benzimidazol-1-yl)acetic acid (triethylammonium salt)

##### A. (7-Nitro-1*H*-benzimidazol-1-yl)acetonitrile

A solution (1 M) of potassium *tert*-butoxide (16.1 ml)) was slowly added to a solution of 4(7)-nitro-1*H*-benzimidazole (2.50 g, 15.3 mmol) in dry DMF (100 ml) at 0-5°C and the resulting dark-red solution was stirred for 15 min at room temperature. Bromoacetonitrile (1.12 mL, 16.1 mmol) was added in one portion and the reaction mixture was stirred for an additional hour, then quenched with dry ice and poured into 400 mL of cold water. The resulting clear solution was repeatedly extracted with  $\text{CHCl}_3$  (4  $\times$  80 ml). Organic extracts were pooled and washed with water (3  $\times$  50 ml) and brine, dried over  $\text{Na}_2\text{SO}_4$  and concentrated, yielding a 1:1 mixture of (4-nitro-1*H*-benzimidazol-1-yl)acetonitrile and (7-nitro-1*H*-benzimidazol-1-yl)acetonitrile. The regioisomers were separated on preparative HPLC (XTerra  $\text{C}_8$  column 19 $\times$ 300 mm, 0.1 M aqueous  $\text{NH}_4\text{Ac}/\text{CH}_3\text{CN}$ ), to yield (7-nitro-1*H*-benzimidazol-1-yl)acetonitrile, 1.15 g (37%). MS (ESI)  $m/z$ : 203.05 [ $\text{M}+\text{H}$ ].  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  ppm 5.68 (s, 2 H) 7.50 (t,  $J=7.8$  Hz, 1 H) 8.16 (m, 1 H) 8.18 (dd,  $J=8.1$ , 1.0 Hz, 1 H) 8.57 (s, 1 H).

30

B. (7-Nitro-1*H*-benzimidazol-1-yl)acetonitrile (1.1 g, 5.4 mmol) was dissolved in 18% hydrochloric acid (30 ml), the solution was transferred into a vial, which was sealed and

heated at 105 °C for 6 h. The vial was cooled, the volatiles were removed under reduced pressure and the residue was co-evaporated two times with acetonitrile. To the residue were added dichloromethane (15 ml) and triethylamine (1 ml), and the slurry was purified on a silica gel column using a mixture of dichloromethane/methanol/triethylamine 84:15:1 (v/v/v) as an eluent to yield the title compound, 1.2 g (69%). MS (ESI)  $m/z$ : 221.98 [M-Et<sub>3</sub>N+H]. <sup>1</sup>H NMR (400 MHz, DMSO-D<sub>6</sub>)  $\delta$  ppm 1.14 (t,  $J$ =7.1 Hz, 9 H) 2.97 (q,  $J$ =7.1 Hz, 6 H) 5.01 (s, 2 H) 7.36 (t,  $J$ =8.1 Hz, 1 H) 7.93 (dd,  $J$ =8.1, 1.0 Hz, 1 H) 8.06 (m, 1 H) 8.37 (s, 1 H).

**2) [7-(Methoxycarbonyl)-1H-benzimidazol-1-yl]acetic acid**

**A. 2-[(2-Hydroxyethyl)amino]-3-nitrobenzoic acid**

2-Chloro-3-nitrobenzoic acid (5.0g, 24.8 mmol) was suspended in ethanol (90 ml) and ethanolamine (4.5 mL, 74.8 mmol) was added. The resulting clear solution was heated at 100°C for two days. The volatiles were removed under reduced pressure. The residue was treated with water (40 ml) and the mixture was acidified with 1M hydrochloric acid to pH 2. A yellow precipitate formed was collected by filtration and washed with water to yield 2-(2-hydroxyethylamino)-3-nitrobenzoic acid, 5.14g (92%). MS (ESI)  $m/z$  225 [M-H]. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD)  $\delta$  ppm 3.04 (t,  $J$ =5.31 Hz, 2 H), 3.69 (t,  $J$ =5.31 Hz, 2 H), 6.71 (t,  $J$ =7.96 Hz, 1 H), 7.93 (dd,  $J$ =8.21, 1.64 Hz, 1 H), 8.13 (dd,  $J$ =7.71, 1.64 Hz, 1 H).

**B. Methyl 2-[(2-hydroxyethyl)amino]-3-nitrobenzoate**

2-(2-Hydroxyethylamino)-3-nitrobenzoic acid (5.14 g, 22.7 mmol) was dissolved in methanol (200 ml) and concentrated H<sub>2</sub>SO<sub>4</sub> (10 ml) was added. The mixture was heated at reflux for 2.5 h. The solvent was removed at reduced pressure. The residue was treated with water (100 ml) and extracted with ethyl acetate (3x150 ml). The combined organic phase was dried and concentrated. Purification by column chromatography on silica using heptane ethyl acetate 1:1 as an eluent afforded methyl 2-[(2-hydroxyethyl)amino]-3-nitrobenzoate, 3.92g (72%). MS (ESI)  $m/z$  241 [M+H]. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  ppm 3.12 (t,  $J$ =5.10 Hz, 2 H), 3.84 (t,  $J$ =5.15 Hz, 2 H), 3.91 (s, 3 H), 6.69 (t,  $J$ =7.96 Hz, 1 H), 7.95 (dd,  $J$ =8.34, 1.52 Hz, 1 H), 8.08 (dd,  $J$ =7.83, 1.52 Hz, 1 H).

**C. Methyl 1-(2-hydroxyethyl)-1H-benzimidazole-7-carboxylate**

Suspension of methyl 2-[(2-hydroxyethyl)amino]-3-nitrobenzoate (3.06 g, 12.7 mmol) in methanol (130 ml) was hydrogenated at atmospheric pressure over 10% palladium on activated charcoal for 10 min. The mixture was filtered through a pad of Celite and the solvent was removed in vacuum. The residue was dissolved in formic acid (60 ml) and heated at 100°C for 45 min and then kept at ambient temperature overnight. Excess of the formic acid was removed under reduced pressure. The residue was dissolved in methanol (100 ml) and treated with concentrated ammonia in methanol (20 ml) for 50 min followed by evaporation of the volatiles. Purification by column chromatography on silica using dichloromethane in methanol 95:5 afforded methyl 1-(2-hydroxyethyl)-1H-benzimidazole-7-carboxylate, 2.31 g (83%). MS (ESI)  $m/z$  221 [M+H]. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD)  $\delta$  ppm 3.78 (t,  $J=5.05$  Hz, 2 H), 3.96 (s, 3 H), 4.70 (t,  $J=5.05$  Hz, 2 H), 7.33 (t,  $J=7.83$  Hz, 1 H), 7.84 - 7.91 (m, 2 H), 8.20 (s, 1 H).

D. To a solution of methyl 1-(2-hydroxyethyl)-1H-benzimidazole-7-carboxylate (2.83 g, 12.8 mmol) in acetone (140 ml) a solution of CrO<sub>3</sub> (1.77 g, 17.7 mmol) and concentrated H<sub>2</sub>SO<sub>4</sub> (1.77 ml) in water (5 ml) was added. The resulting yellow solution was stirred at ambient temperature for 1 h, while the mixture had changed colour to blue green, and then was quenched by the addition of isopropanol. The volatiles were removed in vacuum. The residue was treated with brine and pH of the solution was adjusted to 3 by addition of aqueous sodium bicarbonate. The water phase was repeatedly extracted with ethyl acetate containing 5% methanol. Drying of the organic phase with sodium sulfate, evaporation of solvent and purification of the residue by column chromatography on silica using a gradient of 10-25% methanol in dichloromethane afforded the title compound, 1.44 g (48%). MS (ESI)  $m/z$  235 [M+H]. <sup>1</sup>H NMR (400 MHz, D<sub>2</sub>O)  $\delta$  ppm 3.95 (s, 3 H), 5.17 (s, 2 H), 7.57 (t,  $J=7.95$  Hz, 1 H), 7.96-8.05 (m, 2 H), 8.79 (s, 1 H).

### 3) (7-Cyano-1H-benzimidazol-1-yl)acetic acid

#### A. 2-[(2-Hydroxyethyl)amino]-3-nitrobenzonitrile

A solution of 2-chloro-3-nitrobenzonitrile [prepared as described in WO 97/38983] (0.26 g, 1.4 mmol) and ethanolamine (0.22 mL, 3.5 mmol) in dry ethanol (3.8 ml) was irradiated in a microwave oven at 135 °C for 180 min. The reaction mixture was concentrated under reduced pressure. The residue was dissolved in ethyl acetate, the organic phase was



washed with potassium bisulfate (0.1 M), water and brine, dried over  $\text{Na}_2\text{SO}_4$  and concentrated. Purification was performed using flash chromatography on a silica column and 25% ethyl acetate in heptane as an eluent to yield 2-[(2-hydroxyethyl)amino]-3-nitrobenzonitrile, 0.28 g (95%).  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{CN}$ )  $\delta$  ppm: 3.00 (t,  $J=4.8$  Hz, 1 H), 3.68 (q,  $J=4.7$  Hz, 2 H), 3.81 (m, 2 H), 6.70 (dd,  $J=8.6$ , 7.6 Hz, 1 H), 7.75 (dd,  $J=7.6$ , 1.5 Hz, 1 H), 8.28 (dd,  $J=8.6$ , 2.0 Hz, 1 H), 8.41 (bs, 1 H).

### B. 3-Amino-2-[(2-hydroxyethyl)amino]benzonitrile

To a solution of 2-[(2-hydroxyethyl)amino]-3-nitrobenzonitrile (1.55 g, 7.5 mmol) in a mixture of methanol (30 ml) and water (15 ml) sodium acetate trihydrate (56 g) was added. To this mixture titanium trichloride (65 mL, as 15% solution in 10% aqueous HCl) was added drop-wise over period of 20 min. The resulting dark solution was allowed to stir for additional 2 h, and then carefully neutralized with saturated aqueous sodium bicarbonate. The solids were filtered off, and washed with ethyl acetate. The combined organic phase was washed with water and brine, dried over  $\text{Na}_2\text{SO}_4$  and concentrated yielding 3-amino-2-[(2-hydroxyethyl)amino]benzonitrile (1.23 g, 93%) that was used in the next step without further purification. MS (ESI)  $m/z$ : 178  $[\text{M}+\text{H}]$ .

### C. 1-(2-Hydroxyethyl)-1H-benzimidazole-7-carbonitrile

3-Amino-2-[(2-hydroxyethyl)amino]benzonitrile (1 g, 5.4 mmol) was dissolved in formic acid (3 ml) and irradiated in microwave oven at 135 °C for 2 h. The mixture was cooled and treated with 37% hydrochloric acid (1 ml) at 50°C for 0.5 h. The volatiles were removed under reduced pressure. The residue was partitioned between ethyl acetate and saturated aqueous sodium bicarbonate. The organic phase was washed with water and brine, dried over sodium sulfate and concentrated to yield 1-(2-hydroxyethyl)-1H-benzimidazole-7-carbonitrile, 0.9 g (90%). MS (ESI)  $m/z$  188.1  $[\text{M}+\text{H}]$ .  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  ppm: 3.81 (q,  $J=5.1$  Hz, 2 H), 4.53 (t,  $J=5.3$  Hz, 2 H), 5.03 (t,  $J=5.1$  Hz, 1 H) 7.36 (t,  $J=7.8$  Hz, 1 H), 7.76 (dd,  $J=7.6$ , 1.0 Hz, 1 H), 8.04 (dd,  $J=8.1$ , 1.0 Hz, 1 H), 8.37 (s, 1 H).

D. To a solution of 1-(2-hydroxyethyl)-1H-benzimidazole-7-carbonitrile (0.86 g, 4.6 mmol) in acetone (150 ml) Jones reagent (a mixture of  $\text{CrO}_3$  0.5 g, 5 mmol;  $\text{H}_2\text{SO}_4$  0.5 mL in a minimal amount of water to form a clear solution) was added. The reaction mixture

was stirred for 6 h, quenched with 2-propanol (2 ml) and concentrated to a quarter of the initial volume. The residue was partitioned between ethyl acetate and aqueous potassium hydrosulfate (0.1 M). The aqueous phase was extracted 3-4 times with ethyl acetate and the combined organic extract was washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The oily residue was dissolved in a mixture of dichloromethane (15 ml) and triethylamine (2 ml) and the resulting slurry was loaded onto a flash silica column and eluted with a mixture of dichloromethane/methanol/triethylamine 84:15:1. Fractions containing product were pooled, diluted with dioxane (20 ml), evaporated to dryness and dried *in vacuo* at 40 °C to yield the title product: (7-Cyano-1*H*-benzimidazol-1-yl)acetic acid, 0.36 g (39%). MS (ESI) *m/z* 202.0 [M+H]. <sup>1</sup>H NMR (400 MHz, DMSO-D<sub>6</sub>) δ ppm: 5.31 (s, 2 H), 7.37 (dd, *J*=8.1, 7.7 Hz, 1 H), 7.75 (dd, *J*=7.6, 0.8 Hz, 1 H), 8.04 (dd, *J*=8.1, 1.1 Hz, 1 H), 8.38 (s, 1 H), 13.43 (bs, 1H).

#### 4) (7-Acetyl-1*H*-benzimidazol-1-yl)acetic acid

##### A. 1-[1-(2-Hydroxyethyl)-1*H*-benzimidazol-7-yl]ethanone.

A solution of 1-(2-hydroxyethyl)-1*H*-benzimidazole-7-carbonitrile (0.29 g, 1.5 mmol) in dry THF (6.2 ml) was cooled to -78 °C and MeLi (5.8 mL, 9.3 mmol) was added slowly. After the addition the reaction mixture was allowed to warm up to ambient temperature and kept such for 30 min. Then the temperature was brought down to -78 °C again and water (4 ml) was added slowly. After warming up the reaction mixture was acidified to pH 4 and heated at 50 °C for 30 min. Solvents were removed under reduced pressure and the residue was partitioned between ethyl acetate and aq. NaHCO<sub>3</sub>. The organic extract was further washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. Purification was performed on flash silica column using ethyl acetate – methanol as the eluent.

Yield 0.25 g (80%). Calculated for C<sub>11</sub>H<sub>12</sub>N<sub>2</sub>O<sub>3</sub> *m/z*: 204.23, found 205.23 [M+H]<sup>+</sup>.

<sup>1</sup>H NMR (400 MHz, DMSO-D<sub>6</sub>) δ ppm 2.67 (s, 3 H) 3.51 (q, *J*=5.1 Hz, 2 H) 4.41 (t, *J*=5.3 Hz, 2 H) 4.77 (t, *J*=5.1 Hz, 1 H) 7.29 (t, *J*=7.8 Hz, 1 H) 7.78 (dd, *J*=7.6, 1.0 Hz, 1 H) 7.88 (dd, *J*=8.1, 1.0 Hz, 1 H) 8.20 (s, 1 H).

B. The title compound: (7-acetyl-1*H*-benzimidazol-1-yl)acetic acid, was prepared and isolated as a triethylammonium salt according to the procedure described for the synthesis of (7-Cyano-1*H*-benzimidazol-1-yl)acetic acid (part D). Yield 116 mg (30%). Calculated for

$C_{11}H_{10}N_2O_3$   $m/z$ : 218.21, found 219.16  $[M+H]^+$ .  $^1H$  NMR (400 MHz, DMSO- $D_6$ )  $\delta$  ppm 1.02 (t,  $J=7.1$  Hz, 9 H) 2.56 (s, 3 H) 2.68 - 2.77 (m, 6 H) 4.91 (s, 2 H) 7.24 (t,  $J=7.8$  Hz, 1 H) 7.70 (d,  $J=7.6$  Hz, 1 H) 7.81 - 7.85 (m, 1 H) 8.16 (s, 1 H).

5 **5) (7-Pyridin-2-yl-1H-benzimidazol-1-yl)acetic acid**

**A. Methyl (7-bromo-1H-benzimidazol-1-yl)acetate**

To a solution of (7-bromo-1H-benzimidazol-1-yl)acetic acid triethylamine salt (0.42 g, 1.2 mmol) in methanol (20 ml), conc.  $H_2SO_4$  (2.3 ml) was added and the resulting mixture was heated under reflux for 2 h. After cooling the mixture was concentrated to  $\frac{1}{4}$  of original  
10 volume and partitioned between ethyl acetate and aq.  $NaHCO_3$ . The organic extract was further washed with water and brine, dried over  $Na_2SO_4$  and concentrated.

Yield 0.38 g (97%). Calculated for  $C_{10}H_9BrN_2O_2$   $m/z$ : 267.99, found 269.08  $[M+H]^+$ .

$^1H$  NMR (400 MHz, DMSO- $D_6$ )  $\delta$  ppm 3.72 (s, 3 H) 5.42 (s, 2 H) 7.15 (t,  $J=7.8$  Hz, 1 H) 7.41 - 7.46 (m, 1 H) 7.69 (dd,  $J=8.1$ , 1.0 Hz, 1 H) 8.25 (s, 1 H).

15

**B. Methyl (7-pyridin-2-yl-1H-benzimidazol-1-yl)acetate**

To a mixture of methyl (7-bromo-1H-benzimidazol-1-yl)acetate (108 mg, 0.4 mmol),  $Pd(dppb)Cl_2$  (12 mg), copper(II) oxide (32 mg) in DMF (1.6 ml) under argon, 2-(tributylstannyl)pyridine (0.19 mL, 0.48 mmol) in DMF (0.4 ml) was added in one portion. The  
20 reaction mixture was heated at 100 °C for 23 h in a sealed vial. The vial was cooled and opened and the contents were filtered and concentrated. Purification was performed on flash silica column using heptane – ethyl acetate.

Yield 56 mg (52%). Calculated for  $C_{15}H_{13}N_3O_2$   $m/z$ : 267.10, found 268.12  $[M+H]^+$ .

$^1H$  NMR (400 MHz, MeOD)  $\delta$  ppm 3.42 (s, 3 H) 5.02 (s, 2 H) 7.33 (dd,  $J=7.3$ , 1.3 Hz, 1 H) 7.39 (t,  $J=7.8$  Hz, 1 H) 7.43 - 7.50 (m, 1 H) 7.59 - 7.66 (m, 1 H) 7.79 (dd,  $J=7.8$ , 1.3 Hz, 1 H) 7.91 - 7.99 (m, 1 H) 8.16 (s, 1 H) 8.58 - 8.65 (m, 1 H).

25

**C. (7-Pyridin-2-yl-1H-benzimidazol-1-yl)acetic acid triethylamine salt.**

Methyl (7-pyridin-2-yl-1H-benzimidazol-1-yl)acetate (50 mg, 0.19 mmol) was dissolved in  
30 3 mL methanol and 2 M aq. NaOH (3 mL) was added. The resulting solution was heated at 45 °C until the completion of hydrolysis (3 h) and then concentrated to dryness. The residue was acidified with 5 M aq. HCl, concentrated to dryness, then redissolved in a mixture

of dichloromethane (5 ml) and triethylamine (0.7 ml) and the resulting slurry was loaded onto a flash silica column and eluted with a mixture of dichloromethane/methanol/triethylamine 84:15:1. Fractions containing product were pooled, diluted with dioxane (10 ml), evaporated to dryness and dried under vacuum at 40 °C to yield the title product, 31 mg (47%).

Calculated for  $C_{14}H_{11}N_3O_2$   $m/z$ : 253.09, found 254.14  $[M+H]^+$ .

**6) 5,6-Dihydro-4H-imidazo[4,5,1-ij]quinoline-4-carboxylic acid Hydrochloride**

**A. Methyl 8-amino-1,2,3,4-tetrahydroquinoline-2-carboxylate**

Palladium on carbon (10%, 54 mg) was added to a solution of methyl 4-chloro-8-nitroquinoline-2-carboxylate (127 mg, 0.476 mmol) in ethyl acetate (8 ml) and methanol (8 ml), and the mixture was hydrogenated at 1 atmosphere for 40 min. The catalyst was filtered off, and platinum(IV) oxide (56 mg) was added to the filtrate. The mixture was hydrogenated over 3 h at 1 atmosphere. The catalyst was filtered off, and the filtrate was concentrated *in vacuo*. The residue was purified by column chromatography on silica using heptane/ethyl acetate, 60:40, as the eluent affording 28 mg (29% yield) of the title compound as a yellow oil. MS (ESI)  $m/z$  207  $[M+H]$ .  $^1H$  NMR (400 MHz, DMSO- $D_6$ )  $\delta$  ppm 1.95-2.00 (m, 2 H), 2.50-2.54 (m, 1 H, partly overlapped with the DMSO peak), 2.60-2.67 (m, 1 H), 3.66 (s, 3 H), 4.08-4.11 (m, 1 H), 4.39 (s, 2 H), 4.82 (d,  $J = 2.8$  Hz, 1 H), 6.22 (m, 1 H), 6.33 (t,  $J = 7.4$  Hz, 1 H), 6.38-6.40 (m, 1 H).

**B.** A solution of methyl 8-amino-1,2,3,4-tetrahydroquinoline-2-carboxylate (28 mg, 0.136 mmol) in formic acid (3 ml) was heated at 100 °C for 1 h. The excess of solvent was removed *in vacuo*, and the residual oil was dissolved in a 6 M hydrochloric acid solution and heated at reflux for 30 min. The solvent was removed *in vacuo* affording 32 mg (100% yield) of the title compound as a pink solid. MS (ESI)  $m/z$  203  $[M-HCl+H]$ .  $^1H$  NMR (400 MHz, DMSO- $D_6$ )  $\delta$  ppm 2.36-2.46 (m, 1 H), 2.61-2.66 (m, 1 H), 2.80-2.88 (m, 1 H), 3.10-3.16 (m, 1 H), 5.66 (t,  $J = 4.2$  Hz, 1 H), 7.41 (d,  $J = 7.3$  Hz, 1 H), 7.53 (t,  $J = 7.8$  Hz, 1 H), 7.70 (d,  $J = 8.3$  Hz, 1 H), 9.63 (s, 1 H).

**7) (7-Chloro-6-methoxy-1H-benzimidazol-1-yl)acetic acid**

**A. 2-[(2-Chloro-3-methoxy-6-nitrophenyl)amino]ethanol**

A solution of 2,3-dichloro-1-methoxy-4-nitrobenzene (225 mg, 1.01 mmol) and ethanolamine (309 mg, 5.07 mmol) in ethanol (4 ml) was heated at reflux overnight. Additional ethanolamine (500 mg, 8.20 mmol) was added, and the solution was heated for another 8 h. The solvent was removed *in vacuo*, and the residue was partitioned between water and ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate, and the solvent was removed *in vacuo*. The residue was purified by column chromatography on silica using heptane/ethyl acetate, 70:30, as an eluent affording 141 mg (56% yield) of the title compound as an orange solid. MS (ESI)  $m/z$  247 [M+H].  $^1\text{H}$  NMR (400 MHz, DMSO-D6)  $\delta$  ppm 3.35-3.39 (m, 2 H), 3.50-3.54 (m, 2 H), 3.95 (s, 3 H), 4.85 (t,  $J = 5.0$  Hz, 1 H), 6.75 (d,  $J = 9.6$  Hz, 1 H), 7.10 (broad t,  $J = 5.3$  Hz, 1 H), 8.02 (d,  $J = 9.4$  Hz, 1 H).

#### B. 2-(7-Chloro-6-methoxy-1H-benzimidazol-1-yl)ethanol

The title compound was synthesized according to the procedure described for the synthesis of (7-Cyano-1H-benzimidazol-1-yl)acetic acid, part B and C, starting from 2-[(2-chloro-3-methoxy-6-nitrophenyl)amino]ethanol. Yield 93 mg (74%). MS (ESI)  $m/z$  227 [M+H].  $^1\text{H}$  NMR (400 MHz, DMSO-D6)  $\delta$  ppm 3.72-3.76 (m, 2 H), 3.89 (s, 3 H), 4.51 (t,  $J = 5.6$  Hz, 2 H), 4.95 (t,  $J = 5.3$  Hz, 1 H), 7.10 (d,  $J = 8.8$  Hz, 1 H), 7.58 (d,  $J = 8.6$  Hz, 1 H), 8.05 (s, 1 H).

C. The title compound was synthesized according to the procedure described for the synthesis of (7-Cyano-1H-benzimidazol-1-yl)acetic acid, part D, starting from 2-(7-chloro-6-methoxy-1H-benzimidazol-1-yl)ethanol. Yield 40 mg (44%). MS (ESI)  $m/z$  241 [M+H]. The material was used as such without further purification in the synthesis of the target compound.

#### Syntheses of the intermediates: amines, 8) thru 15)

##### 8) 3-Methoxy-5-(methoxymethyl)aniline

1-Methoxy-3-(methoxymethyl)-5-nitrobenzene (197 mg, 1 mmol) dissolved in methanol (5 ml) was hydrogenated over 10% Pd/C at 40 psi for 2 h at ambient temperature. The reaction mixture was filtered through Celite to remove the catalyst. The filtrate was concentrated in vacuum to yield 3-methoxy-5-(methoxymethyl)aniline (154 mg, 92%).  $^1\text{H}$  NMR

(400 MHz, DMSO-d<sub>6</sub>)  $\delta$  ppm 3.23 (s, 3 H), 3.64 (s, 3H), 4.24 (s, 2 H), 5.01 (br.s, 2 H), 6.41 (br.d, J=7.6 Hz, 1 H), 6.45 (dd, J=8.1, 2.0 Hz, 1 H), 6.51 (t, J=1.7 Hz, 1 H), 6.95 (t, J=8.0 Hz, 1 H)

5    **9) 3-(Methoxymethyl)-5-(trifluoromethyl)aniline**

To a stirred solution of [3-nitro-5-(trifluoromethyl)phenyl]methanol (221 mg, 1 mmol) in THF (1 ml) a solution of potassium tert-butoxide (1M, 1.1 ml, 1.1 mmol) in THF was added at -78°C followed by an addition of methyl iodide (213 mg, 1.5 mmol). The mixture was allowed to reach ambient temperature and was stirred for additional 2 h. The mixture was quenched with water and extracted with chloroform. The extract was dried over sodium sulphate and concentrated in vacuum. The crude product was purified chromatographically on silica gel using 20% ethyl acetate in heptane as an eluent to yield 1-(methoxymethyl)-3-nitro-5-(trifluoromethyl)benzene (130 mg, 55%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  ppm 3.48 (s, 3 H), 4.6 (s, 2 H), 7.93 (s, 1H), 8.38 (br.s, 2 H).

15    1-(Methoxymethyl)-3-nitro-5-(trifluoromethyl)benzene (118 mg, 0.5 mmol) was hydrogenated over 10% Pd/C at 40 psi for 3 h at ambient temperature. The reaction mixture was filtered through Celite to remove the catalyst. The filtrate was concentrated in vacuum to yield 3-(methoxymethyl)-5-(trifluoromethyl)aniline (82 mg, 80%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  ppm 3.39 (s, 3 H), 3.8 (br.s, 2 H), 4.39 (s, 2 H), 6.80 (br.s, 2 H), 6.94 (br.s, 1 H)

20

**10) 1-[3-Amino-5-(trifluoromethyl)phenyl]ethanone**

3-Amino-5-(trifluoromethyl)benzonitrile (186 mg, 1 mmol) dissolved in THF (1 ml) was treated with methyl lithium (1.4M in THF, 2.15 ml, 3 mmol) at -78°C. The mixture was allowed to reach gradually -20°C and stirred for additional 0.5 h. The mixture was quenched with water, acidified with hydrochloric acid to pH 1-2 and warmed gently to 40-45°C for 0.5 h. The mixture was neutralised with sodium bicarbonate and extracted with chloroform. The extract was dried over sodium sulphate and concentrated in vacuum. The crude product was purified using preparative HPLC to yield 1-[3-amino-5-(trifluoromethyl)phenyl]ethanone (108 mg, 53 %). Calculated for C<sub>9</sub>H<sub>8</sub>F<sub>3</sub>NO *m/z*: 203.2, found 204.1 [M+H]<sup>+</sup>. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  ppm 2.58 (s, 3 H), 3.71 (br.s, 2 H), 7.07 (s, 1 H), 7.39 (s, 1 H), 7.52 (s, 1 H)

30

**11) 3-Methoxy-5-(tetrahydrofuran-3-yloxy)aniline****A. 3-(3-Methoxy-5-nitrophenoxy)tetrahydrofuran**

A solution of diethyl azodicarboxylate (40% solution in toluene, 371 mg, 0.85 mmol) in tetrahydrofuran (0.7 ml) was added to a solution of 3-methoxy-5-nitrophenol (111 mg, 0.66 mmol), triphenylphosphine (310 mg, 1.18 mmol), and 3-hydroxytetrahydrofuran (69 mg, 0.79 mmol) in tetrahydrofuran (2 ml). The reaction mixture was stirred at ambient temperature for 4 h. The solvent was removed *in vacuo*, and the residue was partitioned between a 1 M solution of sodium hydroxide and ethyl acetate. The organic layer was washed with a 1 M solution of sodium hydroxide followed by a saturated solution of sodium bicarbonate. The organic layer was dried over magnesium sulfate, and the solvent was removed *in vacuo*. The residue was purified by column chromatography on silica using heptane/ethyl acetate, 90:10→50:50, as an eluent affording 102 mg (65% yield) of the title compound as a pale yellow solid. MS (EI)  $m/z$  239 ( $M^+$ ).  $^1\text{H}$  NMR (400 MHz, DMSO-D<sub>6</sub>)  $\delta$  ppm 1.94-2.01 (m, 1 H), 2.21-2.30 (m, 1 H), 3.74-3.83 (m, 3 H), 3.86 (s, 3 H), 3.87-3.91 (m, 1 H), 5.18-5.21 (m, 1 H), 6.96 (t,  $J$  = 2.3 Hz, 1 H), 7.31 (t,  $J$  = 2.0 Hz, 1 H), 7.34 (t,  $J$  = 2.2 Hz, 1 H).

**B. Palladium on carbon (5%, 30 mg)** was added to a solution of 3-(3-methoxy-5-nitrophenoxy)tetrahydrofuran (100 mg, 0.418 mmol) in ethanol (5 ml) and ethyl acetate (1 ml), and the mixture was hydrogenated at 1 atmosphere for 1 h. The catalyst was filtered off, and the filtrate was concentrated *in vacuo* affording 87 mg (100% yield) of 3-methoxy-5-(tetrahydrofuran-3-yloxy)aniline as an oil. MS (ESI)  $m/z$  210 [ $M+H$ ].  $^1\text{H}$  NMR (400 MHz, DMSO-D<sub>6</sub>)  $\delta$  ppm 1.88-1.95 (m, 1 H), 2.10-2.19 (m, 1 H), 3.62 (s, 3 H), 3.70-3.85 (m, 4 H), 4.83-4.86 (m, 1 H), 5.05 (s, 2 H), 5.64 (t,  $J$  = 2.2 Hz, 1 H), 5.72 (t,  $J$  = 1.9 Hz, 1 H), 5.75 (t,  $J$  = 1.9 Hz, 1 H).

**12) 3-(2-Methoxyethoxy)-5-(trifluoromethyl)aniline****A. 1-(2-Methoxyethoxy)-3-nitro-5-(trifluoromethyl)benzene**

The title compound was synthesized according to the procedure described for the synthesis of 3-methoxy-5-(tetrahydrofuran-3-yloxy)aniline, part A, starting from 3-nitro-5-(trifluoromethyl)phenol and methoxyethanol. Yield 98 mg (50%).  $^1\text{H}$  NMR (400 MHz,

DMSO-D<sub>6</sub>)  $\delta$  ppm 3.32 (s, 3 H, overlapped with water peak), 3.69-3.72 (m, 2 H), 4.35-4.37 (m, 2 H), 7.80 (m, 1 H), 8.03 (t,  $J = 2.2$  Hz, 1 H), 8.05 (m, 1 H).

**B. 3-(2-Methoxyethoxy)-5-(trifluoromethyl)aniline** was synthesized according to the procedure described for the synthesis of 3-methoxy-5-(tetrahydrofuran-3-yloxy)aniline, part B, starting from 1-(2-methoxyethoxy)-3-nitro-5-(trifluoromethyl)benzene. Yield 89 mg. MS (ESI)  $m/z$  236 [M+H]. <sup>1</sup>H NMR (400 MHz, DMSO-D<sub>6</sub>)  $\delta$  ppm 3.30 (s, 3 H), 3.61-3.64 (m, 2 H), 4.03-4.05 (m, 2 H), 5.56 (s, 2 H), 6.31 (s, 1 H), 6.35 (s, 1 H), 6.45 (s, 1 H).

**13) 3-Methoxy-5-(tetrahydrofuran-2-ylmethoxy)aniline**

**A. 2-[(3-Methoxy-5-nitrophenoxy)methyl]tetrahydrofuran**

The title compound was synthesized according to the procedure described for the synthesis of 3-methoxy-5-(tetrahydrofuran-3-yloxy)aniline, part A, starting from 2-(hydroxymethyl)tetrahydrofuran. Yield 104 mg (63%). MS (EI)  $m/z$  253 (M<sup>+</sup>). <sup>1</sup>H NMR (400 MHz, DMSO-D<sub>6</sub>)  $\delta$  ppm 1.64-1.73 (m, 1 H), 1.78-2.04 (m, 3 H), 3.65-3.71 (m, 1 H), 3.76-3.81 (m, 1 H), 3.86 (s, 3 H), 4.00-4.04 (m, 1 H), 4.08-4.12 (m, 1 H), 4.14-4.20 (m, 1 H), 6.98 (t,  $J = 2.3$  Hz, 1 H), 7.32-7.35 (m, 2 H).

**B. 3-Methoxy-5-(tetrahydrofuran-2-ylmethoxy)aniline** was synthesized according to the procedure described for the synthesis of 3-methoxy-5-(tetrahydrofuran-3-yloxy)aniline, part B, starting from 2-[(3-methoxy-5-nitrophenoxy)methyl]tetrahydrofuran. Yield 85 mg (97%). MS (ESI)  $m/z$  224 [M+H]. <sup>1</sup>H NMR (400 MHz, DMSO-D<sub>6</sub>)  $\delta$  ppm 1.58-1.67 (m, 1 H), 1.75-2.01 (m, 3 H), 3.62 (s, 3 H), 3.63-3.68 (m, 1 H), 3.74-3.82 (m, 3 H), 4.06-4.12 (m, 1 H), 5.03 (broad s, 2 H), 5.67 (t,  $J = 2.2$  Hz, 1 H), 5.74 (d,  $J = 2.4$  Hz, 2 H).

**14) 3-methoxy-5-(tetrahydro-2H-pyran-2-ylmethoxy)aniline**

Diisopropyl azodicarboxylate (0.19 mL, 0.99 mmol) was added dropwise to a mixture of *tert*-butyl (3-hydroxy-5-methoxyphenyl)carbamate (196 mg, 0.82 mmol), triphenylphosphine (259 mg, 0.99 mmol), and tetrahydropyran-2-methanol (124 mg, 1.07 mmol) in tetrahydrofuran (2.5 mL) under nitrogen atmosphere. The reaction mixture was stirred at room temperature overnight. The mixture was partitioned between a 1 M NaOH solution and ethyl acetate. The organic layer was washed with brine, dried (MgSO<sub>4</sub>) and evaporated



to give a crude product which was purified by column chromatography to give *tert*-butyl [3-methoxy-5-(tetrahydro-2*H*-pyran-2-ylmethoxy)phenyl]carbamate. This material was treated with 30% solution of trifluoroacetic acid in chloroform overnight. After removal of the volatiles in vacuum 3-methoxy-5-(tetrahydro-2*H*-pyran-2-ylmethoxy)aniline (91 mg, 47%) was isolated as an colourless oil. MS (APCI)  $m/z$  238 [M+H]. <sup>1</sup>H NMR (400 MHz, DMSO-*D*<sub>6</sub>)  $\delta$  ppm 1.22-1.32 (m, 1 H), 1.43-1.51 (m, 3 H), 1.60-1.63 (m, 1 H), 1.79-1.83 (m, 1 H), 3.32-3.40 (m, 1 H), 3.52-3.58 (m, 1 H), 3.61 (s, 3 H), 3.71-3.79 (m, 2 H), 3.86-3.90 (m, 1 H), 5.03 (s, 2 H), 5.66-5.67 (m, 1 H), 5.73-5.74 (m, 2 H).'

### 15 15) 3-(2-isopropoxyethoxy)-5-methoxyaniline

The title compound was synthesized according to the procedure described for the synthesis of 3-methoxy-5-(tetrahydro-2*H*-pyran-2-ylmethoxy)aniline starting from *tert*-butyl (3-hydroxy-5-methoxyphenyl)carbamate and 2-isopropoxyethanol. Yield 78 mg (74%) as an oil. MS (APCI)  $m/z$  226 [M+H]. <sup>1</sup>H NMR (400 MHz, DMSO-*D*<sub>6</sub>)  $\delta$  ppm 1.10 (d,  $J$  = 6.1 Hz, 6 H), 3.58-3.64 (m, 6 H), 3.90-3.92 (m, 2 H), 5.03 (s, 2 H), 5.67 (t,  $J$  = 2.2 Hz, 1 H), 5.74-5.75 (m, 2 H).

## Synthesis of the target compounds

### 20 General method.

To an ice-cooled solution of a 7-substituted (1*H*-benzimidazol-1-yl)acetic acid, prepared as described above (0.14 mmol), triethylamine (0.80 mL, 0.56 mmol) and an appropriate amine (commercially available or described in the literature or described above, 0.2 mmol) in acetonitrile (2 mL) *O*-(7-azabenzotriazol-1-yl)-*N,N,N'*,*N'*-tetramethyluronium hexafluoro-phosphate (69 mg, 0.18 mmol) was added. The ice-bath was removed, and the reaction mixture was stirred at ambient temperature for 0.5 – 3 h. The mixture was quenched with methanol and the volatiles were removed *in vacuo*. The residue was purified by column chromatography on silica using a solution of 0-10% methanol in ethyl acetate as an eluent affording the title compound. Alternatively, the residue was purified by preparative HPLC on XTerra C<sub>8</sub> column (19x300 mm) using 0.1 M aqueous NH<sub>4</sub>OAc/CH<sub>3</sub>CN as an eluent.

Example number	Name	MW calcd	MW found [M+1] or [M-1]	<sup>1</sup> H NMR
1	<i>N</i> -{3-[2-(Dimethyl-amino)ethoxy]phenyl}-2-(7-nitro-1 <i>H</i> -benzimidazol-1-yl)acetamide	383.4	384	(400 MHz, CD <sub>3</sub> OD) δ ppm 2.75 (s, 6 H), 3.30 (t, J=5.1 Hz, 2 H), 4.22 (t, J=5.1 Hz, 2 H), 5.45 (s, 2 H), 6.73 (dd, J=7.6, 2.5 Hz, 1 H), 6.99 (dd, J=8.1, 2.0 Hz, 1 H), 7.23 (t, J=8.1 Hz, 1 H), 7.35 (t, J=2.4 Hz, 1 H), 7.43 (t, J=8.1 Hz, 1 H), 8.04–8.09 (m, 2 H), 8.35 (s, 1 H)
2	<i>N</i> -[3-(Methoxymethyl)phenyl]-2-(7-nitro-1 <i>H</i> -benzimidazol-1-yl)acetamide	340.3	341.1	(400 MHz, CD <sub>3</sub> OD) δ ppm 3.35 (s, 3 H), 4.41 (s, 2 H), 5.45 (s, 2 H), 7.07 (d, J=7.6 Hz, 1 H), 7.28 (t, J=7.9 Hz, 1 H), 7.40–7.51 (m, 3 H), 8.05–8.09 (m, 2 H), 8.35 (s, 1 H)

Example number	Name	MW calcd	MW found [M+1] or [M-1]	<sup>1</sup> H NMR
3	<i>N</i> -(1,3-Dihydro-2-benzofuran-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide	338.3	339.1	(400 MHz, CD <sub>3</sub> OD) δ ppm 5.01 (s, 4 H), 5.40 (s, 2 H), 7.15 (d, J=8.6 Hz, 1 H), 7.29(dd, J=8.1, 1.5 Hz, 1 H), 7.40 (t, J=8.1 Hz, 1 H), 7.47 (d, J=1.5 Hz, 1 H), 8.05 (d, J=8.1 Hz, 1 H), 8.06 (d, J=7.6 Hz, 1 H), 8.24 (s, 1 H)
4	<i>N</i> -[3-Methoxy-5-(methoxymethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide	370.4	371.1	(400 MHz, DMSO-d <sub>6</sub> ) δ ppm 3.30 (s, 3 H), 3.69 (s, 3 H), 4.33 (s, 2 H), 5.38 (s, 2 H), 6.58 (s, 1 H), 7.04 (s, 1 H), 7.10 (t, J=2.1 Hz, 1 H), 7.42 (t, J=7.9 Hz, 1 H), 8.02 (d, J=8.1 Hz, 1 H), 8.14 (dd, J=8.1, 1.1 Hz, 1 H), 8.44 (s, 1 H), 10.39 (br.s, 1 H)
5	<i>N</i> -[3-(Methoxymethyl)-5-(trifluoromethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide	408.3		(400 MHz, CD <sub>3</sub> OD) δ ppm 3.39 (s, 3 H), 4.48 (s, 2 H), 5.47 (s, 2 H), 7.35 (s, 1 H), 7.44 (t, J=8.1 Hz, 1 H), 7.71 (s, 1 H), 7.81 (s, 1 H), 8.08 (d, J=8.1 Hz, 2 H), 8.35 (s, 1 H)

Example number	Name	MW calcd	MW found [M+1] or [M-1]	<sup>1</sup> H NMR
6	<i>N</i> -[3-Cyano-5-(trifluoromethyl)phenyl]-2-(7-nitro-1 <i>H</i> -benzimidazol-1-yl)acetamide	389.3	390.1	(400 MHz, CD <sub>3</sub> OD) δ ppm 5.48 (s, 2 H), 7.42 (t, J=8.1 Hz, 1 H), 7.75 (s, 1 H), 8.07 (d, J=8.1 Hz, 2 H), 8.12 (d, J=7.2 Hz, 2 H), 8.34 (s, 1 H)
7	<i>N</i> -[3-Acetyl-5-(trifluoromethyl)phenyl]-2-(7-nitro-1 <i>H</i> -benzimidazol-1-yl)acetamide	406.3	407.1	(400 MHz, CD <sub>3</sub> OD) δ ppm 2.61 (s, 3 H), 5.51 (s, 2 H), 7.46 (t, J=8.1 Hz, 1 H), 7.95 (s, 1 H), 8.08 - 8.14 (m, 3 H), 8.34 (s, 1 H), 8.36 (s, 1 H)
8	<i>N</i> -[3-(1-Methoxyethyl)phenyl]-2-(7-nitro-1 <i>H</i> -benzimidazol-1-yl)acetamide	354.4	355.1	(400 MHz, CD <sub>3</sub> OD) δ ppm 1.36 (d, J=6.6 Hz, 3 H), 3.18 (s, 3 H), 4.25 (q, J=6.6 Hz, 1 H), 5.40 (d, 2 H), 7.02 (d, J=7.6 Hz, 1 H), 7.25 (t, J=7.8 Hz, 1 H), 7.37 - 7.45 (m, 3 H), 8.05 (d, J=8.1 Hz, 1 H), 8.07 (d, J=8.1 Hz, 1 H), 8.26 (s, 1 H)

Example number	Name	MW calcd	MW found [M+1] or [M-1]	<sup>1</sup> H NMR
9	<i>N</i> -[3-(2-Methoxyethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide	370.4	371	(400 MHz, DMSO-D <sub>6</sub> ) δ ppm 3.29 (s, 3 H), 3.62-3.64 (m, 2 H), 4.01-4.03 (m, 2 H), 5.40 (s, 2 H), 6.65 (dd, <i>J</i> = 8.2, 1.9 Hz, 1 H), 7.04 (d, <i>J</i> = 8.1 Hz, 1 H), 7.19-7.23 (m, 2 H), 7.43 (t, <i>J</i> = 8.0 Hz, 1 H), 8.03 (d, <i>J</i> = 8.1 Hz, 1 H), 8.15 (d, <i>J</i> = 7.6 Hz, 1 H), 8.46 (s, 1 H), 10.39 (s, 1 H)
10	<i>N</i> -[3-Methoxy-5-(2-methoxyethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide	400.4	399	(400 MHz, DMSO-D <sub>6</sub> ) δ ppm 3.29 (s, 3 H), 3.61-3.63 (m, 2 H), 3.69 (s, 3 H), 4.00-4.02 (m, 2 H), 5.38 (s, 2 H), 6.24 (t, <i>J</i> = 2.2 Hz, 1 H), 6.73-6.76 (m, 2 H), 7.43 (t, <i>J</i> = 8.1 Hz, 1 H), 8.03 (dd, 8.1, 0.8 Hz, 1 H), 8.14 (dd, 8.0, 0.9 Hz, 1 H), 8.45 (s, 1 H), 10.36 (s, 1 H)

Example number	Name	MW calcd	MW found [M+1] or [M-1]	<sup>1</sup> H NMR
11	<i>N</i> -[3-(2-Methoxyethoxy)-5-(trifluoromethyl)phenyl]-2-(7-nitro-1 <i>H</i> -benzimidazol-1-yl)acetamide	438.4	437	(400 MHz, DMSO-D <sub>6</sub> ) δ ppm 3.80 (s, 3 H), 3.64-3.66 (m, 2 H), 4.13-4.15 (m, 2 H), 5.43 (s, 2 H), 6.98 (s, 1 H), 7.39 (s, 1 H), 7.44 (t, <i>J</i> = 8.1 Hz, 1 H), 7.50 (s, 1 H), 8.04 (d, <i>J</i> = 7.8 Hz, 1 H), 8.15 (d, <i>J</i> = 8.1 Hz, 1 H), 8.46 (s, 1 H), 10.76 (s, 1 H)
12	<i>N</i> -[3-Methoxy-5-(tetrahydrofuran-2-ylmethoxy)phenyl]-2-(7-nitro-1 <i>H</i> -benzimidazol-1-yl)acetamide	426.4	427	(400 MHz, DMSO-D <sub>6</sub> ) δ ppm 1.61-1.68 (m, 1 H), 1.76-2.01 (m, 3 H), 3.63-3.67 (m, 1 H), 3.69 (s, 3 H), 3.74-3.89 (m, 3 H), 4.08-4.14 (m, 1 H), 5.38 (s, 2 H), 6.24 (t, <i>J</i> = 2.3 Hz, 1 H), 6.72 (t, <i>J</i> = 1.9 Hz, 1 H), 6.77 (t, <i>J</i> = 1.9 Hz, 1 H), 7.43 (t, <i>J</i> = 8.1 Hz, 1 H), 8.03 (dd, <i>J</i> = 8.1, 1.0 Hz, 1 H), 8.14 (dd, <i>J</i> = 8.1, 1.0 Hz, 1 H), 8.45 (s, 1 H), 10.35 (s, 1 H)

Example number	Name	MW calcd	MW found [M+1] or [M-1]	<sup>1</sup> H NMR
13	<i>N</i> -[3-Methoxy-5-(tetrahydrofuran-3-yloxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide	412.4	413	(400 MHz, DMSO-D <sub>6</sub> ) δ ppm 1.90-1.97 (m, 1 H), 2.13-2.22 (m, 1 H), 3.69 (s, 3 H), 3.71-3.86 (m, 4 H), 4.91-4.93 (m, 1 H), 5.38 (m, 2 H), 6.21 (t, <i>J</i> = 2.3 Hz, 1 H), 6.74 (m, 2 H), 7.43 (t, <i>J</i> = 8.1 Hz, 1 H), 8.03 (dd, <i>J</i> = 7.3, 0.8 Hz, 1 H), 8.14 (dd, <i>J</i> = 8.0, 0.9 Hz, 1 H), 8.45 (s, 1 H), 10.37 (s, 1 H)
14	2-(7-Nitro-1H-benzimidazol-1-yl)- <i>N</i> -(3,4,5-trifluorophenyl)acetamide	350.3	351.0	(CD <sub>3</sub> CN) δ ppm 5.31 (s, 2 H) 7.32 (td, <i>J</i> = 10.2, 4.29 Hz, 2 H) 7.41 (t, <i>J</i> = 8.1 Hz, 1 H) 8.05 (dd, <i>J</i> = 8.1, 1.0 Hz, 1 H) 8.08 - 8.11 (m, 2 H) 8.84 (s, 1 H)
15	2-(7-Nitro-1H-benzimidazol-1-yl)- <i>N</i> -(3,4,5-trimethoxybenzyl)acetamide	400.4	401	(400 MHz, DMSO-D <sub>6</sub> ) δ ppm 3.62 (s, 3 H), 3.76 (s, 6 H), 4.22 (d, <i>J</i> = 5.8 Hz, 2 H), 5.24 (s, 2 H), 6.57 (s, 2 H), 7.39 (t, <i>J</i> = 8.0 Hz, 1 H), 7.99 (dd, <i>J</i> = 8.1, 0.8 Hz, 1 H), 8.11 (dd, <i>J</i> = 8.1, 1.0 Hz, 1 H), 8.42 (s, 1 H), 8.72 (t, <i>J</i> = 5.8 Hz, 1 H)

Example number	Name	MW calcd	MW found [M+1] or [M-1]	<sup>1</sup> H NMR
16	<i>N</i> -(3,4-Difluorobenzyl)-2-(7-nitro-1 <i>H</i> -benzimidazol-1-yl)acetamide	346.3	347	(400 MHz, CD <sub>3</sub> OD) δ ppm 4.34 (s, 2 H), 5.32 (s, 2 H), 7.03-7.09 (m, 1 H), 7.12-7.22 (m, 2 H), 7.44 (t, <i>J</i> =8.1 Hz, 1 H), 8.01-8.07 (m, 2 H), 8.32 (s, 1 H)
17	<i>N</i> -[2-(4-Methoxy-phenyl)ethyl]-2-(7-nitro-1 <i>H</i> -benzimidazol-1-yl)acetamide	354.4	355	(400 MHz, CD <sub>3</sub> OD) δ ppm 2.70 (t, <i>J</i> =7.3 Hz, 2 H), 3.35 (t, <i>J</i> =7.3 Hz, 2 H), 3.74 (s, 3 H), 5.22 (s, 2 H), 6.79-6.85 (m, 2 H), 7.09-7.15 (m, 2 H), 7.44 (t, <i>J</i> =8.1 Hz, 1 H), 8.03- 8.09 (m, 2 H), 8.28 (s, 1 H)
18	<i>N</i> -[2-(3-Fluoro-phenyl)ethyl]-2-(7-nitro-1 <i>H</i> -benzimidazol-1-yl)acetamide	342.3	343	(400 MHz, CD <sub>3</sub> OD) δ ppm 2.79 (t, <i>J</i> =7.2 Hz, 2 H), 3.39 (t, <i>J</i> =7.2 Hz, 2 H), 5.25 (s, 2 H), 6.87-6.94 (m, 1 H), 6.96- 7.06 (m, 2 H), 7.24-7.31 (m, 1 H), 7.44 (t, <i>J</i> =8.1 Hz, 1 H), 8.03-8.09 (m, 2 H), 8.29 (s, 1 H)



Example number	Name	MW calcd	MW found [M+1] or [M-1]	<sup>1</sup> H NMR
19	<i>N</i> -[2-(3-Methoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide	355	354.4	(400 MHz, DMF-D9) δ ppm 3.54-3.61 (m, 2 H), 3.80 (s, 3H), 5.36 (s, 2 H), 6.6.78-6.89 (m, 3 H), 7.23 (t, <i>J</i> =7.8 Hz, 1 H), 7.45 (t, <i>J</i> =8.1 Hz, 1 H), 8.05 (dd, <i>J</i> =8.1, 0.8 Hz, 1 H), 8.12 (dd, <i>J</i> =7.8, 1.0 Hz, 1 H), 8.38 (m, 1 H), 8.49 (s, 1 H)
20	2-(7-Nitro-1H-benzimidazol-1-yl)- <i>N</i> -{2-[3-(trifluoromethyl)phenyl]ethyl}acetamide	392.3	393	(400 MHz, DMF-D9) δ ppm 5.35 (s, 2 H), 7.45 (t, <i>J</i> =8.0 Hz, 1 H), 7.58-7.65 (m, 3 H), 7.66 (s, 1 H), 8.05 (dd, <i>J</i> =8.0, 0.9 Hz, 1 H), 8.12 (dd, <i>J</i> =8.1, 1.0 Hz, 1 H) 8.43 (m, 1 H), 8.48 (s, 1 H)
21	<i>N</i> -[2-(3,4-Dimethoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide	384.4	385	(400 MHz, DMF-D9) δ ppm 2.70 (t, <i>J</i> =7.2 Hz, 2 H), 3.37 (q, <i>J</i> =6.7 Hz, 2 H), 3.79 (s, 3 H), 3.82 (s, 3 H), 5.35 (s, 2 H), 6.76 (dd, <i>J</i> =8.2, 1.9 Hz, 1 H), 6.89 (s, 1 H), 6.91 (t, <i>J</i> =1.9 Hz, 1H) 7.45 (t, <i>J</i> =8.1 Hz, 1 H), 8.05 (dd, <i>J</i> =8.1, 0.8 Hz, 1 H), 8.12 (dd, <i>J</i> =8.0, 0.9 Hz, 1 H), 8.36 (m, 1 H), 8.50 (s, 1 H)

Example number	Name	MW calcd	MW found [M+1] or [M-1]	<sup>1</sup> H NMR
22	<i>N</i> -[2-(3,5-Dimethoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide	384.4	385	(400 MHz, CD <sub>3</sub> OD) δ ppm 2.71 (t, <i>J</i> =7.2 Hz, 2 H), 3.39 (t, <i>J</i> =7.2 Hz, 2 H), 3.74 (s, 6 H), 5.25 (s, 2 H), 6.31 (t, <i>J</i> =2.3 Hz, 1 H), 6.41 (d, <i>J</i> =2.2 Hz, 2 H), 7.44 (t, <i>J</i> =8.1 Hz, 1 H), 8.03-8.09 (m, 2 H), 8.29 (s, 1 H)
23	<i>N</i> -(2,3-Dihydro-1H-inden-2-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide	336.4	337	(400 MHz, DMSO-D <sub>6</sub> ) δ ppm 2.72-2.82 (m, 2 H), 3.10-3.19 (m, 2 H), 4.33-4.43 (m, 1 H), 5.13 (s, 2 H), 7.12-7.18 (m, 2 H), 7.20-7.26 (m, 2 H), 7.40 (t, <i>J</i> =8.1 Hz, 1 H), 7.98 (dd, <i>J</i> =8.0, 0.9 Hz, 1 H), 8.11 (dd, <i>J</i> =8.0, 1.0 Hz, 1 H), 8.40 (s, 1 H), 8.57 (d, <i>J</i> =7.1 Hz, 1 H)
24	<i>N</i> -[2-(5-Bromo-2-methoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide	433.3	434	(400 MHz, DMSO-D <sub>6</sub> ) δ ppm 2.65 (t, <i>J</i> =7.1 Hz, 2 H), 3.21 (q, <i>J</i> =6.7 Hz, 2 H), 3.77 (s, 3 H), 5.15 (s, 2 H), 6.94 (d, <i>J</i> =8.6 Hz, 1 H), 7.29-7.43 (m, 3 H), 7.99 (dd, <i>J</i> =8.1, 0.6 Hz, 1 H), 8.10 (dd, <i>J</i> =8.1, 1.0 Hz, 1 H), 8.33 (t, <i>J</i> =5.7 Hz, 1 H), 8.39 (s, 1 H)

Example number	Name	MW calcd	MW found [M+1] or [M-1]	<sup>1</sup> H NMR
25	<i>N</i> -[1-(4-Chlorobenzyl)-2-hydroxyethyl]-2-(7-nitro-1 <i>H</i> -benzimidazol-1-yl)acetamide	388.8	389	(400 MHz, DMSO-D <sub>6</sub> ) δ ppm 2.60-2.68 (m, 1 H), 2.82-2.91 (m, 1 H), 3.25-3.31 (m, 1 H), 3.35-3.43 (m, 1 H), 3.76-3.86 (m, 1 H), 4.83 (t, <i>J</i> =5.4 Hz, 1 H), 5.12 (s, 2 H), 7.18-7.23 (m, 2 H), 7.27-7.33 (m, 2 H), 7.38 (t, <i>J</i> =8.1 Hz, 1 H), 7.97 (dd, <i>J</i> =8.1, 0.8 Hz, 1 H), 8.09 (dd, <i>J</i> =8.0, 0.9 Hz, 1 H), 8.21 (d, <i>J</i> =8.6 Hz, 1 H), 8.38 (s, 1 H)
26	<i>N</i> -(2-Hydroxy-2-phenylethyl)-2-(7-nitro-1 <i>H</i> -benzimidazol-1-yl)acetamide	340.3	341	(400 MHz, DMSO-D <sub>6</sub> ) δ ppm 3.08-3.16 (m, 1 H), 3.20-3.27 (m, 1 H), 4.54 (t, <i>J</i> =6.1 Hz, 1 H), 5.18 (s, 2 H), 5.47 (s, 1 H), 7.20-7.27 (m, 1 H), 7.28-7.35 (m, 4 H), 7.39 (t, <i>J</i> =8.0 Hz, 1 H), 7.98 (dd, <i>J</i> =8.1, 0.8 Hz, 1 H), 8.10 (dd, <i>J</i> =8.1, 1.0 Hz, 1 H), 8.39 (s, 2 H)

Example number	Name	MW calcd	MW found [M+1] or [M-1]	<sup>1</sup> H NMR
27	<i>N</i> -(4-Methoxy-2-naphthyl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide	376.2	377.1	(DMSO-D <sub>6</sub> ) δ ppm 3.93 (s, 3 H) 5.46 (s, 2 H) 7.10 (d, <i>J</i> =1.5 Hz, 1 H) 7.36 (m, 1 H) 7.45 (m, 2 H) 7.71 (m, 2 H) 8.03 (m, 2 H) 8.15 (dd, <i>J</i> =8.1, 1.0 Hz, 1 H) 8.48 (s, 1 H) 10.58 (s, 1 H)
28	2-(7-Acetyl-1H-benzimidazol-1-yl)- <i>N</i> -[3-(methoxymethyl)-5-(trifluoromethyl)phenyl]acetamide	405.4		(400 MHz, CD <sub>3</sub> OD) δ ppm 2.60 (s, 3 H), 3.39 (s, 3 H), 4.48 (s, 2 H), 5.43 (s, 2 H), 7.34 (s, 1 H), 7.39 (t, <i>J</i> =7.8 Hz, 1 H), 7.72 (s, 1 H), 7.85 (s, 1 H), 7.86 (d, <i>J</i> =7.6 Hz, 1 H), 7.92 (d, <i>J</i> =8.1, 1 H), 8.22 (s, 1 H)
29	2-(7-Acetyl-1H-benzimidazol-1-yl)- <i>N</i> -[3-acetyl-5-(trifluoromethyl)phenyl]acetamide	403.4	404.1	(400 MHz, CD <sub>3</sub> OD) δ ppm 2.61 (s, 3 H), 2.62 (s, 3 H), 5.47 (d, 2 H), 7.40 (t, <i>J</i> =8.1 Hz, 1 H), 7.87 – 7.95 (m, 3 H), 8.17 (s, 1 H), 8.23 (s, 1 H), 8.36 (s, 1 H)

Example number	Name	MW calcd	MW found [M+1] or [M-1]	<sup>1</sup> H NMR
30	<i>2-(7-Acetyl-1H-benzimidazol-1-yl)-N-[3-cyano-5-(trifluoromethyl)phenyl]acetamide</i>	386.3	387.1	(400 MHz, CD <sub>3</sub> OD) δ ppm 2.61 (s, 3 H), 5.46 (d, 2 H), 7.39 (t, J=7.9 Hz, 1 H), 7.77 (br.s, 1 H), 7.89 (dd, J=7.6, 1.0 Hz, 1 H), 7.93 (dd, J=8.1, 1.0 Hz, 1 H), 8.14 – 8.18 (m, 2 H), 8.22 (s, 1 H)
31	<i>2-(7-Acetyl-1H-benzimidazol-1-yl)-N-(4-tert-butylbenzyl)acetamide</i>	363.4	364.2	(DMSO-D <sub>6</sub> ) δ ppm 1.23 (s, 9 H) 4.16 (d, J=5.6 Hz, 2 H) 5.12 (s, 2 H) 7.12 - 7.19 (m, J=8.1 Hz, 2 H) 7.26 (t, J=7.8 Hz, 1 H) 7.28 - 7.33 (m, 2 H) 7.72 (dd, J=7.6, 1.0 Hz, 1 H) 7.85 (dd, J=8.1, 1.0 Hz, 1 H) 8.21 (s, 1 H) 8.55 (t, J=5.8 Hz, 1 H)
32	<i>2-(7-Acetyl-1H-benzimidazol-1-yl)-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide</i>	391.3	392.0	(DMSO-D <sub>6</sub> ) δ ppm 2.56 (s, 3 H) 3.79 (s, 3 H) 5.34 (s, 2 H) 6.93 (s, 1 H) 7.32 (t, J=7.8 Hz, 1 H) 7.38 - 7.41 (m, J=2.0 Hz, 1 H) 7.52 (s, 1 H) 7.82 (dd, J=7.6, 1.0 Hz, 1 H) 7.92 (dd, J=8.1, 1.0 Hz, 1 H) 8.27 (s, 1 H) 10.62 (s, 1 H)

Example number	Name	MW calcd	MW found [M+1] or [M-1]	<sup>1</sup> H NMR
33	<i>2-(7-Acetyl-1H-benzimidazol-1-yl)-N-(3,4,5-trimethoxyphenyl)acetamide</i>	383.4	384	(400 MHz, CD <sub>3</sub> OD) δ ppm 2.62 (s, 3 H), 3.72 (s, 3 H), 3.78 (s, 6 H), 5.40 (s, 2 H), 6.90 (s, 2 H), 7.38 (t, <i>J</i> =7.8 Hz, 1 H), 7.86 (dd, <i>J</i> =7.6, 1.0 Hz, 1 H), 7.92 (dd, <i>J</i> =8.1, 1.0 Hz, 1 H), 8.23 (s, 1 H)
34	<i>2-(7-Acetyl-1H-benzimidazol-1-yl)-N-(3,4-difluorophenyl)acetamide</i>	329.3	330	(400 MHz, CD <sub>3</sub> OD) δ ppm 2.61 (s, 3 H), 5.41 (s, 2 H), 7.14-7.24 (m, 2 H), 7.40 (t, <i>J</i> =7.9 Hz, 1 H), 7.56-7.64 (m, 1 H), 7.86 (dd, <i>J</i> =7.7, 0.9 Hz, 1 H), 7.92 (dd, <i>J</i> =8.1, 1.0 Hz, 1 H), 8.22 (s, 1 H)
35	<i>2-(7-Acetyl-1H-benzimidazol-1-yl)-N-(3,5-dimethoxyphenyl)acetamide</i>	353.4	354	(400 MHz, CD <sub>3</sub> OD) δ ppm 2.63 (s, 3 H), 3.74 (s, 6 H), 5.37 (s, 2 H), 6.23 (t, <i>J</i> =2.1 Hz, 1 H), 6.75 (d, <i>J</i> =2.3 Hz, 2 H), 7.38 (t, <i>J</i> =7.8 Hz, 1 H), 7.85 (dd, <i>J</i> =7.6, 0.8 Hz, 1 H), 7.91 (dd, <i>J</i> =8.0, 1.0 Hz, 1 H), 8.21 (s, 1 H)

Example number	Name	MW calcd	MW found [M+1] or [M-1]	<sup>1</sup> H NMR
36	<i>N</i> -(4- <i>tert</i> -Butylbenzyl)-2-(7-cyano-1 <i>H</i> -benzimidazol-1-yl)acetamide	346.4	347	(400 MHz, CD <sub>3</sub> OD) δ ppm 1.29 (s, 9 H), 4.39 (s, 2 H), 5.34 (s, 2 H), 7.24-2.29 (m, 2 H), 7.33-7.38 (m, 2 H), 7.42 (t, <i>J</i> =8.0 Hz, 1 H), 7.71 (dd, <i>J</i> =7.6, 0.8 Hz, 1 H), 8.00 (dd, <i>J</i> =8.1, 1.0 Hz, 1 H), 8.30 (s, 1 H)
37	2-(7-Cyano-1 <i>H</i> -benzimidazol-1-yl)- <i>N</i> -(3,4,5-trimethoxyphenyl)acetamide	366.4	367	(400 MHz, CD <sub>3</sub> OD) δ ppm 3.72 (s, 3 H), 3.79 (s, 6H), 5.47 (s, 2 H), 6.93 (s, 2 H), 7.44 (t, <i>J</i> =8.0 Hz, 1 H), 7.71 (dd, <i>J</i> =7.5, 0.8 Hz, 1 H), 8.02 (dd, <i>J</i> =8.2, 0.9 Hz, 1 H), 8.34 (s, 1 H)
38	<i>N</i> -(4-Bromo-2-fluorophenyl)-2-(7-cyano-1 <i>H</i> -benzimidazol-1-yl)acetamide	373.2	373, 375	(400 MHz, CD <sub>3</sub> OD) δ ppm 5.53 (s, 2 H), 7.28-7.32 (m, 1 H), 7.39-7.45 (m, 2 H), 7.71 (dd, <i>J</i> =7.6, 0.8 Hz, 1 H), 7.91 (t, <i>J</i> =8.6 Hz, 1 H), 8.02 (dd, <i>J</i> =8.4, 1.0 Hz, 1 H), 8.33 (s, 1 H)

Example number	Name	MW calcd	MW found [M+1] or [M-1]	<sup>1</sup> H NMR
39	<i>2-(7-Cyano-1H-benzimidazol-1-yl)-N-(3,4-difluorophenyl)acetamide</i>	312.3	313	(400 MHz, CD <sub>3</sub> OD) δ ppm 5.45 (s, 2 H), 7.16-7.728 (m, 2 H), 7.43 (t, <i>J</i> =8.0 Hz, 1 H), 7.82-7.89 (m, 1 H), 7.71 (dd, <i>J</i> =7.6, 0.8 Hz, 1 H), 8.02 (dd, <i>J</i> =8.2, 0.9 Hz, 1 H), 8.33 (s, 1 H)
40	<i>2-(7-Cyano-1H-benzimidazol-1-yl)-N-(3-ethoxyphenyl)acetamide</i>	320.4	321	(400 MHz, CD <sub>3</sub> OD) δ ppm 1.21 (t, <i>J</i> =7.0 Hz, 3 H), 3.86 (q, <i>J</i> =7.0 Hz, 2 H), 5.32 (s, 2 H), 6.49-6.55 (m, 1 H), 6.88-6.93 (m, 1 H), 7.04 (t, <i>J</i> =8.1 Hz, 1 H), 7.08 (t, <i>J</i> =2.1 Hz, 1 H), 7.29 (t, <i>J</i> =8.0 Hz, 1 H), 7.57 (dd, <i>J</i> =7.5, 0.8 Hz, 1 H), 7.87 (dd, <i>J</i> =8.1, 1.0 Hz, 1 H), 8.18 (s, 1 H)
41	<i>N-(2,3-Dihydro-1H-inden-5-yl)-2-(7-pyridin-2-yl-1H-benzimidazol-1-yl)acetamide</i>	368.4	369.2	(CD <sub>3</sub> CN) δ ppm 1.97 - 2.07 (m, 2 H) 2.77 - 2.85 (m, 4 H) 5.01 (s, 2 H) 6.94 (dd, <i>J</i> =8.1, 1.5 Hz, 1 H) 7.07 (d, <i>J</i> =8.1 Hz, 1 H) 7.12 (s, 1 H) 7.25 - 7.37 (m, 3 H) 7.52 - 7.59 (m, 1 H) 7.68 - 7.76 (m, 1 H) 7.79 (dd, <i>J</i> =7.8, 1.3 Hz, 1 H) 7.98 (s, 1 H) 8.62 - 8.68 (m, 1 H)



Example number	Name	MW calcd	MW found [M+1] or [M-1]	<sup>1</sup> H NMR
42	<i>N</i> -(4- <i>tert</i> -Butylbenzyl)-2-(7-pyridin-2-yl-1 <i>H</i> -benzimidazol-1-yl)acetamide	398.5	399.2	(CD <sub>3</sub> CN) δ ppm 1.21 (s, 9 H) 3.91 (d, <i>J</i> =6.1 Hz, 2 H) 4.82 (s, 2 H) 6.39 (s, 1 H) 6.87 - 6.93 (m, 2 H) 7.16 - 7.27 (m, 5 H) 7.40 - 7.45 (m, Hz, 1 H) 7.64 - 7.71 (m, 2 H) 7.86 (s, 1 H) 8.46 - 8.51 (m, 1 H)
43	2-(7-Chloro-6-methoxy-1 <i>H</i> -benzimidazol-1-yl)- <i>N</i> -(2,3-dihydro-1 <i>H</i> -inden-5-yl)acetamide	355.8	356	(400 MHz, DMSO-D <sub>6</sub> ) δ ppm 1.95-2.02 (m, 2 H), 2.77-2.83 (m, 4 H), 3.87 (s, 3 H), 5.33 (s, 2 H), 7.10 (d, <i>J</i> = 8.8 Hz, 1 H), 7.14 (d, <i>J</i> = 8.1 Hz, 1 H), 7.27 (d, <i>J</i> = 7.8 Hz, 1 H), 7.47 (s, 1 H), 7.60 (d, <i>J</i> = 8.6 Hz, 1 H), 8.13 (s, 1 H), 10.25 (s, 1 H)
44	<i>N</i> -[3-Methoxy-5-(trifluoromethyl)phenyl]-5,6-dihydro-4 <i>H</i> -imidazo[4,5,1- <i>ij</i> ]quinoline-4-carboxamide	375.4	376	(400 MHz, DMSO-D <sub>6</sub> ) δ ppm 2.37-2.44 (m, 1 H), 2.83-3.02 (m, 3 H), 3.81 (s, 3 H), 5.39 (t, <i>J</i> = 4.7 Hz, 1 H), 6.98 (s, 1 H), 7.02 (d, <i>J</i> = 7.1 Hz, 1 H), 7.13 (t, <i>J</i> = 7.6 Hz, 1 H), 7.47-7.49 (m, 2 H), 7.61 (s, 1 H), 8.25 (s, 1 H), 10.76 (s, 1 H)

**Example 45**

2-(7-Amino-1*H*-benzimidazol-1-yl)-*N*-(4-*tert*-butylbenzyl)acetamide

A solution of 2-(7-nitro-1*H*-benzimidazol-1-yl)-*N*-(4-*tert*-butylbenzyl)acetamide (0.35 g, 0.96 mmol) in methanol (15 ml) was hydrogenated in presence of Pd/C catalyst until the consumption of hydrogen ceased. The catalyst was removed by filtration through a pad of Celite™ and concentrated to yield the title compound, 0.32 mg (94%). Calculated for  $C_{20}H_{24}N_4O$  *m/z*: 336.44, found 337.22 [M+H]<sup>+</sup>. <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ ppm 1.25 (s, 9 H) 4.26 (d, *J*=6.1 Hz, 2 H) 5.04 (s, 2 H) 5.06 (s, 2 H) 6.51 (dd, *J*=7.6, 1.0 Hz, 1 H) 6.86 - 6.91 (m, 1 H) 6.91 - 6.95 (m, 1 H) 7.19 (d, *J*=8.1 Hz, 2 H) 7.33 (dt, *J*=8.6, 2.0 Hz, 2 H) 7.93 (s, 1 H) 8.73 (t, *J*=5.6 Hz, 1 H).

#### 10 Example 46

##### *N*-(4-*tert*-Butylbenzyl)-2-(7-iodo-1*H*-benzimidazol-1-yl)acetamide

A suspension of 2-(7-amino-1*H*-benzimidazol-1-yl)-*N*-(4-*tert*-butylbenzyl)acetamide (30 mg, 0.09 mmol) in 2.5M H<sub>2</sub>SO<sub>4</sub> (87 μL) was cooled to 0 °C and 4 M solution of NaNO<sub>2</sub> (25 μL) was added slowly so that the reaction temperature would not exceed 5 °C. After the addition reaction mixture was kept at 0 °C for further 30 min and then added to 1.5 M solution of potassium iodide (100 μL) at ambient temperature. The resulting slurry was partitioned between ethyl acetate and aq. NaHCO<sub>3</sub>. The organic extract was further washed with 1 M Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, water and brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. Purification was performed on flash silica column using ethyl acetate – methanol as the eluent.

15 Yield 18 mg (45%). Calculated for C<sub>20</sub>H<sub>22</sub>IN<sub>3</sub>O<sub>2</sub> *m/z*: 447.31, found 448.06 [M+H]<sup>+</sup>. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>CN) δ ppm 1.29 (s, 9 H) 4.34 (d, *J*=6.1 Hz, 2 H) 5.21 (s, 2 H) 7.00 (t, *J*=7.8 Hz, 1 H) 7.03 - 7.09 (m, 1 H) 7.21 - 7.25 (m, 2 H) 7.37 (dt, *J*=8.6, 2.0 Hz, 2 H) 7.74 (d, *J*=7.6 Hz, 1 H) 7.77 (d, *J*=8.1 Hz, 1 H) 8.11 (s, 1 H).

#### 25 Example 47

##### *N*-(4-*tert*-Butylbenzyl)-2-[7-(dimethylamino)-1*H*-benzimidazol-1-yl]acetamide

To a solution of 2-(7-amino-1*H*-benzimidazol-1-yl)-*N*-(4-*tert*-butylbenzyl)acetamide (24 mg, 66 μmol) and 37% aqueous formaldehyde (100 μL, 1.2 mmol) in ethanol (1 ml), acetic acid (60 μL) and sodium cyanoborohydride (30 mg, 0.5 mmol) were added. After 30 min the volatiles were removed under reduced pressure, and the residue was purified on preparative HPLC to yield the title compound, 15.5 mg (66%). Calculated for C<sub>22</sub>H<sub>28</sub>N<sub>4</sub>O *m/z*: 364.23, found 365.21 [M+H]<sup>+</sup>. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>CN) δ ppm 1.28 (s, 9 H) 2.62 (s, 6

H) 4.30 (d,  $J=6.1$  Hz, 2 H) 5.11 (s, 2 H) 7.07 (m, 2 H) 7.16 (m, 3 H) 7.34 (m, 2 H) 7.41 (dd,  $J=8.1$ , 1.0 Hz, 1 H) 7.87 (s, 1 H).

#### Example 48

5 *2-[7-(1-Hydroxy-1-methylethyl)-1H-benzimidazol-1-yl]-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide*

A solution of 2-(7-acetyl-1H-benzimidazol-1-yl)-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide (26 mg, 0.066 mmol) in dry THF (2.5 ml) was cooled to  $-78^{\circ}\text{C}$ . Methyl magnesium bromide (0.2 mL, 0.2 mmol) was added slowly over a period of 20 min and the reaction was allowed to warm up to  $0^{\circ}\text{C}$  and kept such for additional 1 h. Reaction was quenched with aqueous semi-saturated  $\text{NH}_4\text{Cl}$  and concentrated. The residue was partitioned between ethyl acetate and 0.2 M citric acid (aq.). The organic extract was further washed with  $\text{NaHCO}_3$ , water and brine, dried over  $\text{Na}_2\text{SO}_4$  and concentrated. Purification was performed on reversed-phase preparative HPLC.

15 Yield 15 mg (56%). Calculated for  $\text{C}_{20}\text{H}_{20}\text{F}_3\text{N}_3\text{O}_3$   $m/z$ : 407.39, found 408.03  $[\text{M}+\text{H}]^+$ .

$^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{CN}$ )  $\delta$  ppm 1.57 (s, 6 H) 3.71 (s, 3 H) 5.51 (s, 2 H) 6.84 (bs, 1 H) 7.09 (t,  $J=7.6$  Hz, 1 H) 7.12 - 7.19 (m, 1 H) 7.30 - 7.38 (m, 2 H) 7.53 (dd,  $J=7.8$ , 1.3 Hz, 1 H) 7.84 (s, 1 H) 8.67 (s, 1 H).

#### 20 Example 49

*N-(4-tert-Butylbenzyl)-2-[7-(1-hydroxyethyl)-1H-benzimidazol-1-yl]acetamide*

To a solution of 2-(7-acetyl-1H-benzimidazol-1-yl)-N-(4-tert-butylbenzyl)acetamide (20 mg, 0.054 mmol) in ethanol (3 ml), sodium borohydride (10 mg) was added in single portion. After 30 min the reaction was quenched with acetic acid and concentrated to dryness. The residue was partitioned between ethyl acetate and aq.  $\text{NaHCO}_3$ . The organic extract was further washed with water and brine, dried over  $\text{Na}_2\text{SO}_4$  and concentrated. Purification was performed on flash silica column using ethyl acetate - methanol as the eluent. Yield 20 mg (100%). Calculated for  $\text{C}_{22}\text{H}_{27}\text{N}_3\text{O}_2$   $m/z$ : 365.48, found 366.12  $[\text{M}+\text{H}]^+$ .  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  ppm 1.26 (s, 9 H) 1.40 (d,  $J=6.6$  Hz, 3 H) 4.20 - 4.31 (m, 2 H) 5.05 (m, 1 H) 5.16 - 5.22 (m, 1 H) 5.31 (d,  $J=1.0$  Hz, 1 H) 5.32 - 5.37 (m, 1 H) 7.15 (t,  $J=7.6$  Hz, 1 H) 7.19 (d,  $J=8.6$  Hz, 2 H) 7.27 (d,  $J=7.6$  Hz, 1 H) 7.33 (d,  $J=8.1$  Hz, 2 H) 7.54 (d,  $J=8.1$  Hz, 1 H) 8.10 (s, 1 H) 8.70 (t,  $J=5.8$  Hz, 1 H).

**Example 50***1-{2-[(3,5-Dimethoxyphenyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylic acid*

To [7-(methoxycarbonyl)-1H-benzimidazol-1-yl]acetic acid (0.30 g, 1.28 mmol) in DMF (6 ml) triethylamine (0.89 mL, 6.39 mmol) and 3,5-dimethoxyaniline (0.24 g, 1.54 mmol) were added followed by *O*-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (0.59 g, 1.54 mmol). After stirring the reaction mixture for 1 h the volatiles were removed under reduced pressure. The residue was dissolved in a mixture of THF (10 ml) and water (3 ml) then 10% aqueous NaOH (3 ml) was added. The resulting two-phase reaction mixture was stirred intensively at ambient temperature for 5 h, diluted with water (40 ml) and 1M HCl was added to reach pH 2. Extraction with ethyl acetate : methanol 95:5 (4 x 50 ml), concentration of the combined organic phases and purification of the residue by column chromatography on silica using dichloromethane : methanol 9:1 as eluent afforded the title product as a white solid (0.31 g, 68%). MS (ESI) *m/z*: 354 [M-H]. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ ppm 3.72 (s, 6 H), 5.55 (s, 2 H), 6.22 (t, *J*=2.2 Hz, 1 H), 6.77 (d, *J*=2.0 Hz, 2 H), 7.36 (t, *J*=7.8 Hz, 1 H), 7.89 - 8.00 (m, 2 H), 8.27 (s, 1 H)

**Example 51***1-[2-(2,3-Dihydro-1H-inden-5-ylamino)-2-oxoethyl]-1H-benzimidazole-7-carboxylic acid*

The title compound was synthesized from [7-(methoxycarbonyl)-1H-benzimidazol-1-yl]acetic acid and 2,3-dihydro-1H-inden-5-ylamine according to the procedure described for the preparation of 1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylic acid affording 0.24 g (83%). MS (ESI) *m/z*: 336 [M+H]. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ ppm 2.03 (m, *J*=7.39 Hz, 2 H), 2.79-2.86 (m, 4 H), 5.56 (s, 2 H), 7.09 (d, *J*=8.1 Hz, 1 H), 7.20 (dd, *J*=8.1, 2.0 Hz, 1 H), 7.30 (t, *J*=7.8 Hz, 1 H), 7.38 (s, 1 H), 7.71-7.84 (m, 2 H), 8.20 (s, 1 H)

**Example 52***N-(3,5-Dimethoxyphenyl)-2-[7-(hydroxymethyl)-1H-benzimidazol-1-yl]acetamide*

To 1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylic acid (30 mg, 0.084 mmol) in dry THF (3 ml), 2M BH<sub>3</sub>Me<sub>2</sub>S in THF (0.17 mL, 0.34 mmol) was added keeping the temperature at -20°C to room temperature during a period of 27 h. The

reaction mixture was quenched with acetic acid : water 1:1 (1 ml), the volatiles removed under reduced pressure and the residue purified by preparative HPLC (Xterra C8 column 19x300 mm, 0.1 M aqueous NH<sub>4</sub>Ac/CH<sub>3</sub>CN) giving 1.9 mg (7%) of the desired compound. MS (ESI) m/z: 342 [M+H]. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ ppm 3.73 (s, 6 H), 4.81 (s, 2 H), 5.49 (s, 2 H), 6.25 (t, J=2.3 Hz, 1 H), 6.80 (d, J=2.3 Hz, 2 H), 7.20 - 7.28 (m, 2 H), 7.62-7.68 (m, 1 H), 8.15 (s, 1 H)

### Example 53

#### *1-{2-[(3,5-Dimethoxyphenyl)amino]-2-oxoethyl}-N-ethyl-1H-benzimidazole-7-carboxamide*

To 1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylic acid (20 mg, 0.056 mmol) in DMF (2 ml), triethylamine (39 μL, 0.28 mmol) and *i*-butylchloroformate (8.8 mL, 0.068 mmol) were added. After stirring at room temperature for 10 minutes ethylammonium chloride (5.5 mg, 0.068 mmol) was added, stirring continued for 18 h and the volatiles were removed at reduced pressure. Purification by preparative HPLC (Xterra C8 column 19x300 mm, 0.1 M aqueous NH<sub>4</sub>Ac/CH<sub>3</sub>CN) afforded 13 mg (59%) of the title compound. MS (ESI) m/z: 383 [M+H]. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD), signals given for major (80%) rotamer, δ ppm 0.86 (t, J=7.3 Hz, 3 H), 2.95 (q, J=7.3 Hz, 2 H), 3.77 (s, 6 H), 5.17 (s, 2 H), 6.30 (t, J=2.2 Hz, 1 H), 6.99 (d, J=2.2 Hz, 2 H), 7.36 (t, J=7.8 Hz, 1 H), 7.54 (d, J=7.2 Hz, 1 H), 7.84 (d, J=8.2 Hz, 1 H), 8.20 (s, 1 H)

### Example 54

#### *1-{2-[(3,5-Dimethoxyphenyl)amino]-2-oxoethyl}-N-methyl-1H-benzimidazole-7-carboxamide*

The title compound was prepared according to the procedure described for 1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-N-ethyl-1H-benzimidazole-7-carboxamide starting from 1-{2-[(3,5-Dimethoxyphenyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylic acid and methylammonium chloride affording 14 mg (65%) of the targeted compound. MS (ESI) m/z: 369 [M+H]. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD), signals given for major (75%) rotamer, δ ppm 2.47 (s, 3 H), 3.78 (s, 6 H), 5.17 (s, 2 H), 6.30 (t, J=2.2 Hz, 1 H), 6.98 (d, J=2.2 Hz, 2 H), 7.36 (t, J=7.8 Hz, 1 H), 7.54 (d, J=7.2 Hz, 1 H), 7.84 (dd, J=8.1, 0.9 Hz, 1 H), 8.19 (s, 1 H)

**Example 55**

*1-{2-[(3,5-Dimethoxyphenyl)amino]-2-oxoethyl}-N,N-dimethyl-1H-benzimidazole-7-carboxamide*

- 5 The title compound was prepared according to the procedure described for 1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-N-ethyl-1H-benzimidazole-7-carboxamide starting from 1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylic acid and dimethylammonium chloride affording 6.3 mg (29%) of the targeted compound. MS (ESI) *m/z*: 383 [M+H]. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD), signals given for major (70%)  
10 rotamer, δ ppm 2.63 (s, 3 H), 3.04 (s, 3 H), 3.78 (s, 6 H), 5.40 (s, 2 H), 6.31 (t, *J*=2.3 Hz, 1 H), 6.98 (d, *J*=2.3 Hz, 2 H), 7.36 (t, *J*=7.7 Hz, 1 H), 7.52 (d, *J*=7.3 Hz, 1 H), 7.83 (dd, *J*=8.1, 1.0 Hz, 1 H), 8.15 (s, 1 H)

**Example 56**

- 15 *1-{2-[(3,5-Dimethoxyphenyl)amino]-2-oxoethyl}-N-methoxy-1H-benzimidazole-7-carboxamide*

The title compound was prepared according to the procedure described for 1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-N-ethyl-1H-benzimidazole-7-carboxamide starting from 1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylic acid and methoxyammonium chloride affording 5.5 mg (25%) of the targeted compound. MS (ESI) *m/z*: 385 [M+H]. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ ppm 3.65 (s, 3 H), 3.71 (s, 6 H), 5.42 (s, 2 H), 6.21 (t, *J*=2.3 Hz, 1 H), 6.77 (d, *J*=2.3 Hz, 2 H), 7.33 (t, *J*=7.7 Hz, 1 H), 7.39-7.44 (m, 1 H), 7.86 (dd, *J*=8.1, 1.1 Hz, 1 H), 8.23 (s, 1 H)

- 25 **Example 57**

*Ethyl 1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylate*

- To 1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylic acid (20 mg, 0.056 mmol) in DMF (2 ml) triethylamine (39 μL, 0.28 mmol) and O-benzotriazol-1-yl-N,N,N',N'-tetramethyluronium hexafluorophosphate (26 mg, 0.067 mmol) were  
30 added. The resulting solution was stirred at ambient temperature for 20 minutes followed by addition of ethanol and stirring for additional 20 h. The volatiles were evaporated under reduced pressure and the residue was purified by preparative HPLC (Xterra C8 column

19x300 mm, 0.1 M aqueous NH<sub>4</sub>Ac/CH<sub>3</sub>CN) affording the desired product, 4.5 mg (21%).  
MS (ESI) m/z: 384 [M+H]. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ ppm 1.06 (t, J=7.1 Hz, 3 H),  
3.78 (s, 6 H), 3.98 (q, J=7.1 Hz, 2 H), 5.32 (s, 2 H), 6.30 (t, J=2.3 Hz, 1 H), 6.97 (d, J=2.0  
Hz, 2 H), 7.38 (t, J=7.8 Hz, 1 H), 7.58 (d, J=7.3 Hz, 1 H), 7.85 (d, J=8.3 Hz, 1 H), 8.20 (s,  
1 H)

### Example 58

#### *Ethyl 1-[2-[(4-tert-butylbenzyl)amino]-2-oxoethyl]-1H-benzimidazole-7-carboxylate*

The title compound was prepared according to the procedure described for the preparation  
of ethyl 1-[2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl]-1H-benzimidazole-7-carboxylate  
using 1-[2-[(4-tert-butylbenzyl)amino]-2-oxoethyl]-1H-benzimidazole-7-carboxylic acid  
as starting material which afforded 2.2 mg (5%) of the product. MS (ESI) m/z: 394 [M+H].  
<sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ ppm 1.29 (s, 9 H), 1.34 (t, J=7.2 Hz, 3 H), 4.26 (q, J=7.2  
Hz, 2 H), 4.32 (s, 2 H), 5.41 (s, 2 H), 7.18-7.24 (m, 2 H), 7.31 - 7.37 (m, 3 H), 7.86-7.92  
(m, 2 H), 8.20 (s, 1 H)

### Example 59

#### *Ethyl 1-[2-(2,3-dihydro-1H-inden-5-ylamino)-2-oxoethyl]-1H-benzimidazole-7-carboxylate*

The title product was prepared according to the procedure described for the preparation of  
ethyl 1-[2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl]-1H-benzimidazole-7-carboxylate  
using 1-[2-(2,3-dihydro-1H-inden-5-ylamino)-2-oxoethyl]-1H-benzimidazole-7-carboxylic  
acid as starting material which afforded 6.0 mg (15%) of the product. MS (ESI) m/z: 364  
[M+H]. <sup>1</sup>H NMR (600 MHz, CD<sub>3</sub>OD) δ ppm 1.29 (t, J=7.1 Hz, 3 H), 2.06 (m, 2 H), 2.86  
(m, 4 H), 4.29 (q, J=7.1 Hz, 2 H), 5.52 (s, 2 H), 7.12 (d, J=7.9 Hz, 1 H), 7.21 (d, J=8.1 Hz,  
1 H), 7.36 (t, J=7.8 Hz, 1 H), 7.40 (s, 1 H), 7.87 (d, J=7.5 Hz, 1 H), 7.93 (d, J=8.0 Hz, 1  
H), 8.25 (s, 1 H)

### Example 60

#### *2-(1H-Benzimidazol-1-yl)-N-(4-tert-butylbenzyl)acetamide*

The title product was prepared according to the procedure used for the preparation of com-  
pounds described in examples 1 thru 44. Calculated for C<sub>20</sub>H<sub>23</sub>N<sub>3</sub>O m/z: 321.2, found 322.2

[M+H]<sup>+</sup>. <sup>1</sup>H NMR (400 MHz, DMSO-D<sub>6</sub>) δ ppm 1.26 (s, 9 H) 4.26 (d, *J*=6.1 Hz, 2 H) 4.97 (s, 2 H) 7.16 - 7.27 (m, 4 H) 7.30 - 7.37 (m, 2 H) 7.45 (dd, *J*=7.1, 1.5 Hz, 1 H) 7.61 - 7.68 (m, 1 H) 8.17 (s, 1 H) 8.75 (t, *J*=5.8 Hz, 1 H)

#### 5 Example 61

##### *2-(1H-Benzimidazol-1-yl)-N-(2,3-dihydro-1H-inden-5-yl)acetamide*

The title product was prepared according to the procedure used for the preparation of compounds described in examples 1 thru 44. Calculated for C<sub>20</sub>H<sub>23</sub>N<sub>3</sub>O *m/z*: 291.4, found 292 [M+H]<sup>+</sup>. <sup>1</sup>H NMR (400 MHz, DMSO-D<sub>6</sub>) δ ppm 1.92 - 2.03 (m, *J*=7.4, 2 H), 2.79 (q, *J*=7.3 Hz, 4 H), 5.13 (s, 2 H), 7.14 (d, *J*=8.1 Hz, 1 H), 7.17 - 7.30 (m, 3 H), 7.47 - 7.54 (m, 2 H), 7.63-7.68 (m, 1 H), 8.21 (s, 1 H), 10.32 (s, 1 H)

#### Example 62

##### *N-[3-Methoxy-5-(tetrahydro-2H-pyran-2-ylmethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide*

Synthesised according to the general method of synthesis of the target compounds from (7-Nitro-1H-benzimidazol-1-yl)acetic acid and 3-methoxy-5-(tetrahydro-2H-pyran-2-ylmethoxy)aniline. MS (ESI) *m/z* 441 [M+H]. <sup>1</sup>H NMR (400 MHz, DMSO-D<sub>6</sub>) δ ppm 1.26-1.32 (m, 1 H), 1.45-1.50 (m, 3 H), 1.60-1.62 (m, 1 H), 1.79-1.81 (m, 1 H), 3.35-3.40 (m, partly overlapped with water peak, 1 H), 3.55-3.60 (m, 1 H), 3.69 (s, 3 H), 3.82-3.88 (m, 3 H), 5.38 (s, 2 H), 6.23 (s, 1 H), 6.73 (s, 1 H), 6.75 (s, 1 H), 7.43 (t, *J* = 7.8 Hz, 1 H), 8.03 (d, *J* = 7.6 Hz, 1 H), 8.14 (d, *J* = 7.6 Hz, 1 H), 8.45 (s, 1 H), 10.34 (s, 1 H).

#### Example 63

##### *N-[3-(2-Isopropoxyethoxy)-5-methoxyphenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide*

Synthesised according to the general method of synthesis of the target compounds from (7-Nitro-1H-benzimidazol-1-yl)acetic acid and 3-(2-isopropoxyethoxy)-5-methoxyaniline. MS (ESI) *m/z* 429 [M+H]. <sup>1</sup>H NMR (400 MHz, DMSO-D<sub>6</sub>) δ ppm 1.09 (d, *J* = 6.1 Hz, 6 H), 3.57-3.64 (m, 1 H), 3.64-3.66 (m, 2 H), 3.69 (s, 3 H), 3.97-3.99 (m, 2 H), 5.38 (s, 2 H), 6.24 (t, *J* = 2.2 Hz, 1 H), 6.73 (t, *J* = 1.8 Hz, 1 H), 6.76 (t, *J* = 1.8 Hz, 1 H), 7.43 (t, *J* = 8.1 Hz, 1 H), 8.03 (d, *J* = 7.8 Hz, 1 H), 8.14 (dd, *J* = 7.8, 0.8 Hz, 1 H), 8.45 (s, 1 H), 10.36 (s, 1 H).



**Example 64**

*N*-{3-methoxy-5-[2-(2-oxopyrrolidin-1-yl)ethoxy]phenyl}-2-(7-nitro-1H-benzimidazol-1-yl)acetamide

5

Synthesised according to the general method of synthesis of the target compounds from (7-Nitro-1H-benzimidazol-1-yl)acetic acid and 1-[2-(3-amino-5-methoxy-phenoxy)ethyl]pyrrolidin-2-one. MS (ESI) *m/z* 454 (M+H). <sup>1</sup>H NMR (400 MHz, DMSO-D6) δ ppm 1.86-1.94 (m, 2 H), 2.20 (t, *J* = 8.1 Hz, 2 H), 3.42 (t, *J* = 7.1 Hz, 2 H), 3.51 (t, *J* = 5.4 Hz, 2 H), 3.70 (s, 3 H), 4.00 (t, *J* = 5.6 Hz, 2 H), 5.38 (s, 2 H), 6.25 (t, *J* = 2.2 Hz, 1 H), 6.73 (t, *J* = 1.8 Hz, 1 H), 6.78 (t, *J* = 1.8 Hz, 1 H), 7.43 (t, *J* = 8.1 Hz, 1 H), 8.03 (d, *J* = 8.1 Hz, 1 H), 8.14 (d, *J* = 7.6 Hz, 1 H), 8.45 (s, 1 H), 10.37 (s, 1 H).

10

**Example 65**

*N*-{3-[2-(1H-imidazol-1-yl)ethoxy]-5-methoxyphenyl}-2-(7-nitro-1H-benzimidazol-1-yl)acetamide

15

Synthesised according to the general method of synthesis of the target compounds from (7-Nitro-1H-benzimidazol-1-yl)acetic acid and 3-[2-(1H-imidazol-1-yl)ethoxy]-5-methoxyaniline. MS (ESI) *m/z* 437 (M+H). <sup>1</sup>H NMR (400 MHz, DMSO-D6) δ ppm 3.69 (s, 3 H), 4.16 (t, *J* = 5.1 Hz, 2 H), 4.32 (t, *J* = 5.2 Hz, 2 H), 5.38 (s, 2 H), 6.23 (t, *J* = 2.2 Hz, 1 H), 6.72-6.74 (m, 1 H), 6.77-6.78 (m, 1 H), 6.87 (s, 1 H), 7.21 (s, 1 H), 7.43 (t, *J* = 8.0 Hz, 1 H), 7.65 (s, 1 H), 8.03 (d, *J* = 7.8 Hz, 1 H), 8.14 (dd, *J* = 7.8, 0.8 Hz, 1 H), 8.45 (s, 1 H), 10.37 (s, 1 H).

20

**Pharmacology**

25

**1. hVR1 FLIPR (Fluorometric Image Plate Reader) screening assay**

Transfected CHO cells, stably expressing hVR1 (15,000 cells/well) are seeded in 50 ul media in a black clear bottom 384 plate (Greiner) and grown in a humidified incubator (37°C, 2% CO<sub>2</sub>), 24-30 hours prior to experiment.

30

Subsequently, the media is removed from the cell plate by inversion and 2  $\mu$ M Fluo-4 is added using a multidrop (Labsystems). Following the 40 minutes dye incubation in the dark at 37°C and 2% CO<sub>2</sub>, the extracellular dye present is washed away using an EMBLA (Scatron), leaving the cells in 40ul of assay buffer (1 X HBSS, 10 mM D-Glucose, 1 mM CaCl<sub>2</sub>, 10 mM HEPES, 10 X 7.5% NaHCO<sub>3</sub> and 2.5 mM Probenecid).

*FLIPR assay - IC<sub>50</sub> determination protocol*

For IC<sub>50</sub> determinations the fluorescence is read using FLIPR filter 1 (em 520-545 nm). A cellular baseline recording is taken for 30 seconds, followed by a 20  $\mu$ l addition of 10, titrated half-log concentrations of the test compound, yielding cellular concentration ranging from 3  $\mu$ M to 0.1 nM. Data is collected every 2 seconds for a further 5 minutes prior to the addition of a VR1 agonist solution: either 50 nM solution of capsaicin or MES (2-[N-morpholino] ethanesulfonic acid) buffer (pH 5.2), by the FLIPR pipettor. The FLIPR continues to collect data for a further 4 minutes. Compounds having antagonistic properties against the hVR1 will inhibit the increase in intracellular calcium in response to the capsaicin addition. This consequently leading to a reduction in fluorescence signal and providing a reduced fluorescence reading, compared with no compound, buffer controls. Data is exported by the FLIPR program as a sum of fluorescence calculated under the curve upon the addition of capsaicin. Maximum inhibition, Hill slope and IC<sub>50</sub> data for each compound are generated.

2. DRGs were dissected out from adult Sprague Dawley rats (100-300 gr), and placed on ice in L15 Leibovitz medium. The ganglia were enzyme treated with Collagenase 80U/ml+ Dispace 34 U/ml dissolved in DMEM +5% serum, overnight at 37 °C. The next day, cells were triturated with fire polished pasteur pipettes, and seeded in the center of 58 mm diameter Nunc cell dishes coated with Poly-D Lysine (1 mg/ml). The DRGs were cultured in a defined medium without foetal bovine serum, containing Dulbecco's MEM / NUT MIX F-12 (1:1) without L-glutamine but with pyridoxine, 6 mg/mL D(+)-Glucose, 100  $\mu$ g/mL apo-transferrin, 1 mg/mL BSA, 20  $\mu$ g/mL insulin, 2 mM L-glutamine, 50 IU/ mL Penicillin, 50  $\mu$ g / mL Streptomycin and 0.01  $\mu$ g/mL NGF-7S.

When the cells had grown for 2 days up to 4 weeks, the experiments were done. Cells were chosen based on size and presence of neurites. Small cells with long processes were used for recording (most likely to be C neurons, with native VR1 receptors).

The cells were recorded with conventional whole cell voltage clamp patch clamp, using the following solutions (calcium ion free):

The extracellular solution comprised (in mM): NaCl 137, KCl 5,  $MgCl_2 \cdot H_2O$  1.2, HEPES 10, Glucose 10, EGTA 5, Sucrose 50, pH to 7.4 with NaOH.

The intracellular solution comprised K-gluconate 140, NaCl 3,  $MgCl_2 \cdot H_2O$  1.2, HEPES 10, EGTA 1, pH to 7.2 with KOH. When the cells were penetrated with suction, a puff of capsaicin (500 nM) was used to determine if the cell expressed VR1 receptor. If not, a new cell was chosen. If yes, then the compounds were added in increasing doses before the capsaicin pulse (500 nM), to determine an  $IC_{50}$  value.

#### List of abbreviations

VR1	vanilloid receptor 1
IBS	irritable bowel syndrome
IBD	inflammatory bowel disease
GERD	gastro-esophageal reflux disease
DRG	Dorsal Root Ganglion
BSA	Bovine Serum Albumin
HEPES	4-(2-Hydroxyethyl)piperazine-1-ethanesulfonic acid
EGTA	Ethylene glycol-bis(2-aminoethylether)- <i>N,N,N',N'</i> -tetraacetic acid
DMEM	Dulbeccos Modified Eagle's Medium

#### Results


Typical  $IC_{50}$  values as measured in the assays described above are 10  $\mu M$  or less. In one aspect of the invention the  $IC_{50}$  is below 500 nM. In another aspect of the invention the  $IC_{50}$  is below 100 nM. In a further aspect of the invention the  $IC_{50}$  is below 10 nM.

Table 1. Specimen results from the hVR1 FLIPR.

Example No.	Name	IC <sub>50</sub> nM (agonist)
10	<i>N</i> -[3-Methoxy-5-(2-methoxyethoxy)phenyl]-2-(7-nitro-1 <i>H</i> -benzimidazol-1-yl)acetamide	22 (capsaicin) 45 (low pH)
14	2-(7-nitro-1 <i>H</i> -benzimidazol-1-yl)- <i>N</i> -(3,4,5-trifluorophenyl)acetamide	48 (capsaicin) 108 (low pH)
32	2-(7-acetyl-1 <i>H</i> -benzimidazol-1-yl)- <i>N</i> -[3-methoxy-5-(trifluoromethyl)phenyl]acetamide	77 (capsaicin) 53 (low pH)
35	2-(7-Acetyl-1 <i>H</i> -benzimidazol-1-yl)- <i>N</i> -(3,5-dimethoxyphenyl)acetamide	518 (capsaicin) 508 (low pH)

## CLAIMS

1. A compound having the formula



(I)

wherein:

$R^1$  is H,  $NO_2$ , halo,  $NR^6R^7$ ,  $C_{1-6}alkyl$ ,  $C_{2-6}alkenyl$ ,  $C_{2-6}alkynyl$ ,  $C_{1-6}haloalkyl$ ,  $C_{1-6}haloalkylo$ ,  $R^6OC_{0-6}alkyl$ ,  $R^6CO$ ,  $R^6OCO$  or  $CONR^6R^7$ ;

$m$  is 0, 1, 2 or 3;

$R^2$  is H,  $NO_2$ , halo,  $NR^6R^7$ ,  $C_{1-6}alkyl$ ,  $C_{2-6}alkenyl$ ,  $C_{2-6}alkynyl$ ,  $C_{1-6}haloalkyl$ ,  $C_{1-6}haloalkylo$ , cyano,  $R^6OC_{0-6}alkyl$ ,  $R^6CO$ ,  $R^6OCO$ ,  $R^6CONR^7$ ,  $R^6R^7NCO$ ,  $R^8SO_2$ ,  $R^8SO_2HN$ ,  $arylC_{0-6}alkyl$  or heteroaryl $C_{0-6}alkyl$ ;

$R^3$  and  $R^3$  are each independently H or  $C_{1-4}alkyl$ ;

$R^2$  and  $R^3$  optionally form a ring;

$p$  is 0, 1 or 2;

$n$  is 0, 2, 3 or 4;

$R^5$  is  $C_{1-10}alkyl$ ,  $C_{6-10}arylC_{0-6}alkyl$ ,  $C_{3-7}cycloalkylC_{0-6}alkyl$  or  $C_{5-6}heteroarylC_{0-6}alkyl$ , whereby any aryl, heteroaryl or cycloalkyl may be fused with aryl, heteroaryl,  $C_{3-7}cycloalkyl$  or  $C_{3-7}heterocycloalkyl$ , and which  $R^5$  may be substituted with one or more

A;

A is H, OH,  $NO_2$ , cyano,  $R^6CO$ ,  $R^6O(CO)$ , halo,  $C_{1-6}alkyl$ ,  $NR^6R^7$ ,  $C_{1-6}haloalkyl$ ,  $C_{1-6}haloalkylo$ ,  $R^6OC_{0-6}alkyl$ , hydroxy $C_{1-6}alkyl$ ,  $R^8SO_2$ ,  $R^8SO_2HN$ ,  $C_{5-6}arylo$  or  $CONR^6R^7$ ;

$R^6$  and  $R^7$  are each independently H or  $C_{1-6}alkyl$ ; and

$R^8$  is  $NR^6R^7$  or  $C_{1-4}alkyl$ ,

which compound is selected from the group consisting of

N-[3-[2-(dimethylamino)ethoxy]phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,

- N-[3-(methoxymethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-(1,3-dihydro-2-benzofuran-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-methoxy-5-(methoxymethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-(methoxymethyl)-5-(trifluoromethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-[3-(methoxymethyl)-5-(trifluoromethyl)phenyl]acetamide,  
N-[3-cyano-5-(trifluoromethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-acetyl-5-(trifluoromethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-[3-acetyl-5-(trifluoromethyl)phenyl]acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-[3-cyano-5-(trifluoromethyl)phenyl]acetamide,  
N-[3-(1-methoxyethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-chloro-6-methoxy-1H-benzimidazol-1-yl)-N-(2,3-dihydro-1H-inden-5-yl)acetamide,  
N-[3-(2-methoxyethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-methoxy-5-(2-methoxyethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-(2-methoxyethoxy)-5-(trifluoromethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-methoxy-5-(tetrahydrofuran-2-ylmethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-methoxy-5-(tetrahydrofuran-3-yloxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-methoxy-5-(trifluoromethyl)phenyl]-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-4-carboxamide,  
2-(7-amino-1H-benzimidazol-1-yl)-N-(4-tert-butylbenzyl)acetamide,  
N-(4-tert-butylbenzyl)-2-(7-iodo-1H-benzimidazol-1-yl)acetamide,  
2-[7-(1-hydroxy-1-methylethyl)-1H-benzimidazol-1-yl]-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,  
N-(4-tert-butylbenzyl)-2-[7-(1-hydroxyethyl)-1H-benzimidazol-1-yl]acetamide,  
N-(2,3-dihydro-1H-inden-5-yl)-2-(7-pyridin-2-yl-1H-benzimidazol-1-yl)acetamide,  
N-(4-tert-butylbenzyl)-2-(7-pyridin-2-yl-1H-benzimidazol-1-yl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-(4-tert-butylbenzyl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,

- 2-(7-nitro-1H-benzimidazol-1-yl)-N-(3,4,5-trifluorophenyl)acetamide,  
N-(4-tert-butylbenzyl)-2-(7-cyano-1H-benzimidazol-1-yl)acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-(3,4,5-trimethoxyphenyl)acetamide,  
N-(4-bromo-2-fluorophenyl)-2-(7-cyano-1H-benzimidazol-1-yl)acetamide,  
5 2-(7-cyano-1H-benzimidazol-1-yl)-N-(3,4-difluorophenyl)acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-(3-ethoxyphenyl)acetamide,  
2-(7-nitro-1H-benzimidazol-1-yl)-N-(3,4,5-trimethoxybenzyl)acetamide,  
N-(3,4-difluorobenzyl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[2-(4-methoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
10 N-[2-(3-fluorophenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[2-(3-methoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-nitro-1H-benzimidazol-1-yl)-N-{2-[3-(trifluoromethyl)phenyl]ethyl}acetamide,  
N-[2-(3,4-dimethoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-(3,4,5-trimethoxyphenyl)acetamide,  
15 2-(7-acetyl-1H-benzimidazol-1-yl)-N-(3,4-difluorophenyl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-(3,5-dimethoxyphenyl)acetamide,  
N-[2-(3,5-dimethoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-(2,3-dihydro-1H-inden-2-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[2-(5-bromo-2-methoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
20 N-[1-(4-chlorobenzyl)-2-hydroxyethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-(2-hydroxy-2-phenylethyl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylic acid,  
1-[2-(2,3-dihydro-1H-inden-5-ylamino)-2-oxoethyl]-1H-benzimidazole-7-carboxylic acid,  
N-(3,5-dimethoxyphenyl)-2-[7-(hydroxymethyl)-1H-benzimidazol-1-yl]acetamide,  
25 1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-N-ethyl-1H-benzimidazole-7-carbox-  
amide,  
1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-N-methyl-1H-benzimidazole-7-carbox-  
amide,  
1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-N,N-dimethyl-1H-benzimidazole-7-car-  
30 boxamide,  
1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-N-methoxy-1H-benzimidazole-7-car-  
boxamide,

- ethyl 1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylate,  
ethyl 1-{2-[(4-tert-butylbenzyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylate,  
ethyl 1-[2-(2,3-dihydro-1H-inden-5-ylamino)-2-oxoethyl]-1H-benzimidazole-7-carboxy-  
late,
- 5 N-(4-tert-butylbenzyl)-2-[7-(dimethylamino)-1H-benzimidazol-1-yl]acetamide,  
N-(4-methoxy-2-naphthyl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(1H-benzimidazol-1-yl)-N-(4-tert-butylbenzyl)acetamide,  
2-(1H-benzimidazol-1-yl)-N-(2,3-dihydro-1H-inden-5-yl)acetamide,  
N-(3,5-dimethoxyphenyl)-2-(7-ethynyl-1H-benzimidazol-1-yl)acetamide,
- 10 N-(3,5-dimethoxyphenyl)-2-(7-prop-1-yn-1-yl-1H-benzimidazol-1-yl)acetamide,  
N-(3,5-dimethoxyphenyl)-2-[7-(1H-1,2,3-triazol-4-yl)-1H-benzimidazol-1-yl]acetamide,,  
N-(3,5-dimethoxyphenyl)-2-[7-(1-methyl-1H-1,2,3-triazol-4-yl)-1H-benzimidazol-1-  
yl]acetamide,  
N-(3,5-dimethoxyphenyl)-2-[7-(1-methyl-1H-tetrazol-5-yl)-1H-benzimidazol-1-
- 15 yl]acetamide,  
2-(7-ethyl-1H-benzimidazol-1-yl)-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,  
2-[7-(2-hydroxyethyl)-1H-benzimidazol-1-yl]-N-[3-methoxy-5-(trifluoro-  
methyl)phenyl]acetamide,  
2-[7-(2-hydroxy-1-methylethyl)-1H-benzimidazol-1-yl]-N-[3-methoxy-5-(trifluoro-
- 20 methyl)phenyl]acetamide,  
N-[3-methoxy-5-(trifluoromethyl)phenyl]-2-(7-vinyl-1H-benzimidazol-1-yl)acetamide,  
2-(7-isopropenyl-1H-benzimidazol-1-yl)-N-[3-methoxy-5-(trifluoro-  
methyl)phenyl]acetamide ,  
2-(7-isopropyl-1H-benzimidazol-1-yl)-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,
- 25 N-(3,5-dimethoxyphenyl)-2-(7-methoxy-1H-benzimidazol-1-yl)acetamide,  
N-(3,5-dimethoxyphenyl)-2-(7-ethoxy-1H-benzimidazol-1-yl)acetamide,  
N-(3,5-dimethoxyphenyl)-2-(7-isopropoxy-1H-benzimidazol-1-yl)acetamide,  
2-(7-tert-butoxy-1H-benzimidazol-1-yl)-N-(3,5-dimethoxyphenyl)acetamide,  
N-(3,5-dimethoxyphenyl)-2-[7-(trifluoromethoxy)-1H-benzimidazol-1-yl]acetamide
- 30 N-(3,5-dimethoxyphenyl)-2-[7-(methylsulfinyl)-1H-benzimidazol-1-yl]acetamide,  
2-[7-(difluoromethyl)-1H-benzimidazol-1-yl]-N-(3,5-dimethoxyphenyl)acetamide,  
2-[7-(cyanomethyl)-1H-benzimidazol-1-yl]-N-(3,5-dimethoxyphenyl)acetamide,



- 2-[7-(aminomethyl)-1H-benzimidazol-1-yl]-N-(3,5-dimethoxyphenyl)acetamide  
N-(3,5-dimethoxyphenyl)-2-{7-[(dimethylamino)methyl]-1H-benzimidazol-1-yl}acetamide,  
2-(7-cyclopropyl-1H-benzimidazol-1-yl)-N-(3,5-dimethoxyphenyl)acetamide,  
5 2-(7-cyclobutyl-1H-benzimidazol-1-yl)-N-(3,5-dimethoxyphenyl)acetamide,  
N-(3,5-dimethoxyphenyl)-2-[7-(methoxymethyl)-1H-benzimidazol-1-yl]acetamide,  
N-(1-isopropyl-1H-benzimidazol-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-fluoro-1H-benzimidazol-1-yl)-N-(1-isopropyl-1H-benzimidazol-5-yl)acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-(1-isopropyl-1H-benzimidazol-5-yl)acetamide,  
10 N-(1-tert-butyl-1H-benzimidazol-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-(1-tert-butyl-2-methyl-1H-benzimidazol-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-(1-isopropyl-2-methyl-1H-benzimidazol-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
15 2-(7-cyano-1H-benzimidazol-1-yl)-N-(1-isopropyl-2-methyl-1H-benzimidazol-5-yl)acetamide,  
N-(1-tert-butyl-2-methyl-1H-benzimidazol-5-yl)-2-(7-cyano-1H-benzimidazol-1-yl)acetamide,  
N-(1-tert-butyl-1H-benzimidazol-5-yl)-2-(7-cyano-1H-benzimidazol-1-yl)acetamide,  
20 N-(1-tert-butyl-1H-benzimidazol-5-yl)-2-(7-fluoro-1H-benzimidazol-1-yl)acetamide,  
N-(1-tert-butyl-2-methyl-1H-benzimidazol-5-yl)-2-(7-fluoro-1H-benzimidazol-1-yl)acetamide,  
2-(7-fluoro-1H-benzimidazol-1-yl)-N-(1-isopropyl-2-methyl-1H-benzimidazol-5-yl)acetamide,  
25 N-[1-isopropyl-7-(trifluoromethyl)-1H-benzimidazol-5-yl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-fluoro-1H-benzimidazol-1-yl)-N-[1-isopropyl-7-(trifluoromethyl)-1H-benzimidazol-5-yl]acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-[1-isopropyl-7-(trifluoromethyl)-1H-benzimidazol-5-yl]acetamide,  
30 5-yl]acetamide,  
N-2-naphthyl-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-2-naphthylacetamide,

- 2-(7-fluoro-1H-benzimidazol-1-yl)-N-2-naphthylacetamide,  
 2-(7-cyano-1H-benzimidazol-1-yl)-N-(4-methoxy-2-naphthyl)acetamide,  
 2-(7-fluoro-1H-benzimidazol-1-yl)-N-(4-methoxy-2-naphthyl)acetamide,  
 N-[3-methoxy-5-(tetrahydro-2H-pyran-2-ylmethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-  
 1-yl)acetamide,  
 N-[3-(2-isopropoxyethoxy)-5-methoxyphenyl]-2-(7-nitro-1H-benzimidazol-1-  
 yl)acetamide,  
 N-[3,5-bis(2-ethoxyethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
 N-[3-methoxy-5-[2-(2-oxopyrrolidin-1-yl)ethoxy]phenyl]-2-(7-nitro-1H-benzimidazol-1-  
 yl)acetamide,  
 N-[3-methoxy-5-(3-morpholin-4-ylpropoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-  
 yl)acetamide,  
 N,N-diethyl-2-(3-methoxy-5-[(7-nitro-1H-benzimidazol-1-  
 yl)acetyl]amino}phenoxy)acetamide,  
 N-[3-methoxy-5-[(1-methylpiperidin-2-yl)methoxy]phenyl]-2-(7-nitro-1H-benzimidazol-  
 1-yl)acetamide,  
 N-[3-[2-(1H-imidazol-1-yl)ethoxy]-5-methoxyphenyl]-2-(7-nitro-1H-benzimidazol-1-  
 yl)acetamide, and  
 N-[3-methoxy-5-[(1-methyl-1H-imidazol-5-yl)methoxy]phenyl]-2-(7-nitro-1H-benzimi-  
 dazol-1-yl)acetamide,  
 or salts, solvates or solvated salts thereof.

2. A compound according to claim 1 selected from the group consisting of  
 N-[3-[2-(dimethylamino)ethoxy]phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
 N-[3-(methoxymethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
 N-(1,3-dihydro-2-benzofuran-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
 N-[3-methoxy-5-(methoxymethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
 N-[3-(methoxymethyl)-5-(trifluoromethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-  
 yl)acetamide,  
 2-(7-acetyl-1H-benzimidazol-1-yl)-N-[3-(methoxymethyl)-5-(trifluoro-  
 methyl)phenyl]acetamide,  
 N-[3-cyano-5-(trifluoromethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,

- N-[3-acetyl-5-(trifluoromethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-[3-acetyl-5-(trifluoromethyl)phenyl]acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-[3-cyano-5-(trifluoromethyl)phenyl]acetamide,  
N-[3-(1-methoxyethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-chloro-6-methoxy-1H-benzimidazol-1-yl)-N-(2,3-dihydro-1H-inden-5-yl)acetamide,  
N-[3-(2-methoxyethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-methoxy-5-(2-methoxyethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-(2-methoxyethoxy)-5-(trifluoromethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-methoxy-5-(tetrahydrofuran-2-ylmethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-methoxy-5-(tetrahydrofuran-3-yloxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-methoxy-5-(trifluoromethyl)phenyl]-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-4-carboxamide,  
2-(7-amino-1H-benzimidazol-1-yl)-N-(4-tert-butylbenzyl)acetamide,  
N-(4-tert-butylbenzyl)-2-(7-iodo-1H-benzimidazol-1-yl)acetamide,  
2-[7-(1-hydroxy-1-methylethyl)-1H-benzimidazol-1-yl]-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,  
N-(4-tert-butylbenzyl)-2-[7-(1-hydroxyethyl)-1H-benzimidazol-1-yl]acetamide,  
N-(2,3-dihydro-1H-inden-5-yl)-2-(7-pyridin-2-yl-1H-benzimidazol-1-yl)acetamide,  
N-(4-tert-butylbenzyl)-2-(7-pyridin-2-yl-1H-benzimidazol-1-yl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-(4-tert-butylbenzyl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,  
2-(7-nitro-1H-benzimidazol-1-yl)-N-(3,4,5-trifluorophenyl)acetamide,  
N-(4-tert-butylbenzyl)-2-(7-cyano-1H-benzimidazol-1-yl)acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-(3,4,5-trimethoxyphenyl)acetamide,  
N-(4-bromo-2-fluorophenyl)-2-(7-cyano-1H-benzimidazol-1-yl)acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-(3,4-difluorophenyl)acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-(3-ethoxyphenyl)acetamide,  
2-(7-nitro-1H-benzimidazol-1-yl)-N-(3,4,5-trimethoxybenzyl)acetamide,  
N-(3,4-difluorobenzyl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,

- N-[2-(4-methoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[2-(3-fluorophenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[2-(3-methoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-nitro-1H-benzimidazol-1-yl)-N-{2-[3-(trifluoromethyl)phenyl]ethyl} acetamide,  
5 N-[2-(3,4-dimethoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-(3,4,5-trimethoxyphenyl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-(3,4-difluorophenyl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-(3,5-dimethoxyphenyl)acetamide,  
N-[2-(3,5-dimethoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
10 N-(2,3-dihydro-1H-inden-2-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[2-(5-bromo-2-methoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[1-(4-chlorobenzyl)-2-hydroxyethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-(2-hydroxy-2-phenylethyl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylic acid,  
15 1-[2-(2,3-dihydro-1H-inden-5-ylamino)-2-oxoethyl]-1H-benzimidazole-7-carboxylic acid,  
N-(3,5-dimethoxyphenyl)-2-[7-(hydroxymethyl)-1H-benzimidazol-1-yl]acetamide,  
1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-N-ethyl-1H-benzimidazole-7-carbox-  
amide,  
1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-N-methyl-1H-benzimidazole-7-carbox-  
20 amide,  
1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-N,N-dimethyl-1H-benzimidazole-7-car-  
boxamide,  
1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-N-methoxy-1H-benzimidazole-7-car-  
boxamide,  
25 ethyl 1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylate,  
ethyl 1-{2-[(4-tert-butylbenzyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylate,  
ethyl 1-[2-(2,3-dihydro-1H-inden-5-ylamino)-2-oxoethyl]-1H-benzimidazole-7-carboxy-  
late,  
N-(4-tert-butylbenzyl)-2-[7-(dimethylamino)-1H-benzimidazol-1-yl]acetamide,  
30 N-(4-methoxy-2-naphthyl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(1H-benzimidazol-1-yl)-N-(4-tert-butylbenzyl)acetamide,  
2-(1H-benzimidazol-1-yl)-N-(2,3-dihydro-1H-inden-5-yl)acetamide,

N-(3,5-dimethoxyphenyl)-2-(7-ethynyl-1H-benzimidazol-1-yl)acetamide,  
N-(3,5-dimethoxyphenyl)-2-(7-prop-1-yn-1-yl-1H-benzimidazol-1-yl)acetamide,  
N-(3,5-dimethoxyphenyl)-2-[7-(1H-1,2,3-triazol-4-yl)-1H-benzimidazol-1-yl]acetamide,,  
N-(3,5-dimethoxyphenyl)-2-[7-(1-methyl-1H-1,2,3-triazol-4-yl)-1H-benzimidazol-1-  
5 yl]acetamide,  
N-(3,5-dimethoxyphenyl)-2-[7-(1-methyl-1H-tetrazol-5-yl)-1H-benzimidazol-1-  
yl]acetamide,  
2-(7-ethyl-1H-benzimidazol-1-yl)-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,  
2-[7-(2-hydroxyethyl)-1H-benzimidazol-1-yl]-N-[3-methoxy-5-(trifluoro-  
10 methyl)phenyl]acetamide,  
2-[7-(2-hydroxy-1-methylethyl)-1H-benzimidazol-1-yl]-N-[3-methoxy-5-(trifluoro-  
methyl)phenyl]acetamide,  
N-[3-methoxy-5-(trifluoromethyl)phenyl]-2-(7-vinyl-1H-benzimidazol-1-yl)acetamide,  
2-(7-isopropenyl-1H-benzimidazol-1-yl)-N-[3-methoxy-5-(trifluoro-  
15 methyl)phenyl]acetamide ,  
2-(7-isopropyl-1H-benzimidazol-1-yl)-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,  
N-(3,5-dimethoxyphenyl)-2-(7-methoxy-1H-benzimidazol-1-yl)acetamide,  
N-(3,5-dimethoxyphenyl)-2-(7-ethoxy-1H-benzimidazol-1-yl)acetamide,  
N-(3,5-dimethoxyphenyl)-2-(7-isopropoxy-1H-benzimidazol-1-yl)acetamide,  
20 2-(7-tert-butoxy-1H-benzimidazol-1-yl)-N-(3,5-dimethoxyphenyl)acetamide,  
N-(3,5-dimethoxyphenyl)-2-[7-(trifluoromethoxy)-1H-benzimidazol-1-yl]acetamide  
N-(3,5-dimethoxyphenyl)-2-[7-(methylsulfinyl)-1H-benzimidazol-1-yl]acetamide,  
2-[7-(difluoromethyl)-1H-benzimidazol-1-yl]-N-(3,5-dimethoxyphenyl)acetamide,  
2-[7-(cyanomethyl)-1H-benzimidazol-1-yl]-N-(3,5-dimethoxyphenyl)acetamide,  
25 2-[7-(aminomethyl)-1H-benzimidazol-1-yl]-N-(3,5-dimethoxyphenyl)acetamide  
N-(3,5-dimethoxyphenyl)-2-{7-[(dimethylamino)methyl]-1H-benzimidazol-1-  
yl}acetamide,  
2-(7-cyclopropyl-1H-benzimidazol-1-yl)-N-(3,5-dimethoxyphenyl)acetamide,  
2-(7-cyclobutyl-1H-benzimidazol-1-yl)-N-(3,5-dimethoxyphenyl)acetamide,  
30 N-(3,5-dimethoxyphenyl)-2-[7-(methoxymethyl)-1H-benzimidazol-1-yl]acetamide,  
N-(1-isopropyl-1H-benzimidazol-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-fluoro-1H-benzimidazol-1-yl)-N-(1-isopropyl-1H-benzimidazol-5-yl)acetamide,

- 2-(7-cyano-1H-benzimidazol-1-yl)-N-(1-isopropyl-1H-benzimidazol-5-yl)acetamide,  
N-(1-tert-butyl-1H-benzimidazol-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-(1-tert-butyl-2-methyl-1H-benzimidazol-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
5 N-(1-isopropyl-2-methyl-1H-benzimidazol-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-(1-isopropyl-2-methyl-1H-benzimidazol-5-yl)acetamide,  
N-(1-tert-butyl-2-methyl-1H-benzimidazol-5-yl)-2-(7-cyano-1H-benzimidazol-1-yl)acetamide,  
10 N-(1-tert-butyl-1H-benzimidazol-5-yl)-2-(7-cyano-1H-benzimidazol-1-yl)acetamide,  
N-(1-tert-butyl-1H-benzimidazol-5-yl)-2-(7-fluoro-1H-benzimidazol-1-yl)acetamide,  
N-(1-tert-butyl-2-methyl-1H-benzimidazol-5-yl)-2-(7-fluoro-1H-benzimidazol-1-yl)acetamide,  
15 2-(7-fluoro-1H-benzimidazol-1-yl)-N-(1-isopropyl-2-methyl-1H-benzimidazol-5-yl)acetamide,  
N-[1-isopropyl-7-(trifluoromethyl)-1H-benzimidazol-5-yl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-fluoro-1H-benzimidazol-1-yl)-N-[1-isopropyl-7-(trifluoromethyl)-1H-benzimidazol-5-yl]acetamide,  
20 5-yl]acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-[1-isopropyl-7-(trifluoromethyl)-1H-benzimidazol-5-yl]acetamide,  
N-2-naphthyl-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-2-naphthylacetamide,  
25 2-(7-fluoro-1H-benzimidazol-1-yl)-N-2-naphthylacetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-(4-methoxy-2-naphthyl)acetamide,  
2-(7-fluoro-1H-benzimidazol-1-yl)-N-(4-methoxy-2-naphthyl)acetamide,  
N-[3-methoxy-5-(tetrahydro-2H-pyran-2-ylmethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
30 N-[3-(2-isopropoxyethoxy)-5-methoxyphenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide, and

N-[3-methoxy-5-(3-morpholin-4-ylpropoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
or salts, solvates or solvated salts thereof.

3. A compound according to claim 1 selected from the group consisting of
- N-{3-[2-(dimethylamino)ethoxy]phenyl}-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-(methoxymethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-(1,3-dihydro-2-benzofuran-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-methoxy-5-(methoxymethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,
- 10 N-[3-(methoxymethyl)-5-(trifluoromethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-[3-(methoxymethyl)-5-(trifluoromethyl)phenyl]acetamide,  
N-[3-cyano-5-(trifluoromethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,
- 15 N-[3-acetyl-5-(trifluoromethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-[3-acetyl-5-(trifluoromethyl)phenyl]acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-[3-cyano-5-(trifluoromethyl)phenyl]acetamide,  
N-[3-(1-methoxyethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-chloro-6-methoxy-1H-benzimidazol-1-yl)-N-(2,3-dihydro-1H-inden-5-yl)acetamide,
- 20 N-[3-(2-methoxyethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-methoxy-5-(2-methoxyethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-(2-methoxyethoxy)-5-(trifluoromethyl)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-methoxy-5-(tetrahydrofuran-2-ylmethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,
- 25 y)acetamide,  
N-[3-methoxy-5-(tetrahydrofuran-3-yloxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-methoxy-5-(trifluoromethyl)phenyl]-5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-4-carboxamide,
- 30 2-(7-amino-1H-benzimidazol-1-yl)-N-(4-tert-butylbenzyl)acetamide,  
N-(4-tert-butylbenzyl)-2-(7-iodo-1H-benzimidazol-1-yl)acetamide,

- 2-[7-(1-hydroxy-1-methylethyl)-1H-benzimidazol-1-yl]-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,  
N-(4-tert-butylbenzyl)-2-[7-(1-hydroxyethyl)-1H-benzimidazol-1-yl]acetamide,  
N-(2,3-dihydro-1H-inden-5-yl)-2-(7-pyridin-2-yl-1H-benzimidazol-1-yl)acetamide,  
5 N-(4-tert-butylbenzyl)-2-(7-pyridin-2-yl-1H-benzimidazol-1-yl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-(4-tert-butylbenzyl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,  
2-(7-nitro-1H-benzimidazol-1-yl)-N-(3,4,5-trifluorophenyl)acetamide,  
N-(4-tert-butylbenzyl)-2-(7-cyano-1H-benzimidazol-1-yl)acetamide,  
10 2-(7-cyano-1H-benzimidazol-1-yl)-N-(3,4,5-trimethoxyphenyl)acetamide,  
N-(4-bromo-2-fluorophenyl)-2-(7-cyano-1H-benzimidazol-1-yl)acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-(3,4-difluorophenyl)acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-(3-ethoxyphenyl)acetamide,  
2-(7-nitro-1H-benzimidazol-1-yl)-N-(3,4,5-trimethoxybenzyl)acetamide,  
15 N-(3,4-difluorobenzyl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[2-(4-methoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[2-(3-fluorophenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[2-(3-methoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-nitro-1H-benzimidazol-1-yl)-N-{2-[3-(trifluoromethyl)phenyl]ethyl}acetamide,  
20 N-[2-(3,4-dimethoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-(3,4,5-trimethoxyphenyl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-(3,4-difluorophenyl)acetamide,  
2-(7-acetyl-1H-benzimidazol-1-yl)-N-(3,5-dimethoxyphenyl)acetamide,  
N-[2-(3,5-dimethoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
25 N-(2,3-dihydro-1H-inden-2-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[2-(5-bromo-2-methoxyphenyl)ethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[1-(4-chlorobenzyl)-2-hydroxyethyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-(2-hydroxy-2-phenylethyl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
1-[2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl]-1H-benzimidazole-7-carboxylic acid,  
30 1-[2-(2,3-dihydro-1H-inden-5-ylamino)-2-oxoethyl]-1H-benzimidazole-7-carboxylic acid,  
N-(3,5-dimethoxyphenyl)-2-[7-(hydroxymethyl)-1H-benzimidazol-1-yl]acetamide,



1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-N-ethyl-1H-benzimidazole-7-carboxamide,

1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-N-methyl-1H-benzimidazole-7-carboxamide,

5 1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-N,N-dimethyl-1H-benzimidazole-7-carboxamide,

1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-N-methoxy-1H-benzimidazole-7-carboxamide,

ethyl 1-{2-[(3,5-dimethoxyphenyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylate,

10 ethyl 1-{2-[(4-tert-butylbenzyl)amino]-2-oxoethyl}-1H-benzimidazole-7-carboxylate,

ethyl 1-{2-[(2,3-dihydro-1H-inden-5-ylamino)-2-oxoethyl]-1H-benzimidazole-7-carboxylate,

N-(4-tert-butylbenzyl)-2-[7-(dimethylamino)-1H-benzimidazol-1-yl]acetamide,

N-(4-methoxy-2-naphthyl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,

15 2-(1H-benzimidazol-1-yl)-N-(4-tert-butylbenzyl)acetamide, and

2-(1H-benzimidazol-1-yl)-N-(2,3-dihydro-1H-inden-5-yl)acetamide,

or salts, solvates or solvated salts thereof.

4. A compound according to claim 1 selected from the group consisting of

20 N-(3,5-dimethoxyphenyl)-2-(7-ethynyl-1H-benzimidazol-1-yl)acetamide,

N-(3,5-dimethoxyphenyl)-2-(7-prop-1-yn-1-yl-1H-benzimidazol-1-yl)acetamide,

N-(3,5-dimethoxyphenyl)-2-[7-(1H-1,2,3-triazol-4-yl)-1H-benzimidazol-1-yl]acetamide,,

N-(3,5-dimethoxyphenyl)-2-[7-(1-methyl-1H-1,2,3-triazol-4-yl)-1H-benzimidazol-1-yl]acetamide,

25 N-(3,5-dimethoxyphenyl)-2-[7-(1-methyl-1H-tetrazol-5-yl)-1H-benzimidazol-1-yl]acetamide,

2-(7-ethyl-1H-benzimidazol-1-yl)-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,

2-[7-(2-hydroxyethyl)-1H-benzimidazol-1-yl]-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,

30 2-[7-(2-hydroxy-1-methylethyl)-1H-benzimidazol-1-yl]-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,

N-[3-methoxy-5-(trifluoromethyl)phenyl]-2-(7-vinyl-1H-benzimidazol-1-yl)acetamide,

2-(7-isopropenyl-1H-benzimidazol-1-yl)-N-[3-methoxy-5-(trifluoro-methyl)phenyl]acetamide,

2-(7-isopropyl-1H-benzimidazol-1-yl)-N-[3-methoxy-5-(trifluoromethyl)phenyl]acetamide,  
N-(3,5-dimethoxyphenyl)-2-(7-methoxy-1H-benzimidazol-1-yl)acetamide,

- 5 N-(3,5-dimethoxyphenyl)-2-(7-ethoxy-1H-benzimidazol-1-yl)acetamide,  
N-(3,5-dimethoxyphenyl)-2-(7-isopropoxy-1H-benzimidazol-1-yl)acetamide,  
2-(7-tert-butoxy-1H-benzimidazol-1-yl)-N-(3,5-dimethoxyphenyl)acetamide,  
N-(3,5-dimethoxyphenyl)-2-[7-(trifluoromethoxy)-1H-benzimidazol-1-yl]acetamide  
N-(3,5-dimethoxyphenyl)-2-[7-(methylsulfinyl)-1H-benzimidazol-1-yl]acetamide,  
10 2-[7-(difluoromethyl)-1H-benzimidazol-1-yl]-N-(3,5-dimethoxyphenyl)acetamide,  
2-[7-(cyanomethyl)-1H-benzimidazol-1-yl]-N-(3,5-dimethoxyphenyl)acetamide,  
2-[7-(aminomethyl)-1H-benzimidazol-1-yl]-N-(3,5-dimethoxyphenyl)acetamide  
N-(3,5-dimethoxyphenyl)-2-{7-[(dimethylamino)methyl]-1H-benzimidazol-1-yl}acetamide,

- 15 2-(7-cyclopropyl-1H-benzimidazol-1-yl)-N-(3,5-dimethoxyphenyl)acetamide,  
2-(7-cyclobutyl-1H-benzimidazol-1-yl)-N-(3,5-dimethoxyphenyl)acetamide,  
N-(3,5-dimethoxyphenyl)-2-[7-(methoxymethyl)-1H-benzimidazol-1-yl]acetamide,  
N-(1-isopropyl-1H-benzimidazol-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-fluoro-1H-benzimidazol-1-yl)-N-(1-isopropyl-1H-benzimidazol-5-yl)acetamide,  
20 2-(7-cyano-1H-benzimidazol-1-yl)-N-(1-isopropyl-1H-benzimidazol-5-yl)acetamide,  
N-(1-tert-butyl-1H-benzimidazol-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-(1-tert-butyl-2-methyl-1H-benzimidazol-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-(1-isopropyl-2-methyl-1H-benzimidazol-5-yl)-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
25 y)acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-(1-isopropyl-2-methyl-1H-benzimidazol-5-yl)acetamide,  
N-(1-tert-butyl-2-methyl-1H-benzimidazol-5-yl)-2-(7-cyano-1H-benzimidazol-1-yl)acetamide,  
30 N-(1-tert-butyl-1H-benzimidazol-5-yl)-2-(7-cyano-1H-benzimidazol-1-yl)acetamide,  
N-(1-tert-butyl-1H-benzimidazol-5-yl)-2-(7-fluoro-1H-benzimidazol-1-yl)acetamide,

- N-(1-tert-butyl-2-methyl-1H-benzimidazol-5-yl)-2-(7-fluoro-1H-benzimidazol-1-yl)acetamide,  
2-(7-fluoro-1H-benzimidazol-1-yl)-N-(1-isopropyl-2-methyl-1H-benzimidazol-5-yl)acetamide,  
5 N-[1-isopropyl-7-(trifluoromethyl)-1H-benzimidazol-5-yl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-fluoro-1H-benzimidazol-1-yl)-N-[1-isopropyl-7-(trifluoromethyl)-1H-benzimidazol-5-yl]acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-[1-isopropyl-7-(trifluoromethyl)-1H-benzimidazol-10 5-yl]acetamide,  
N-2-naphthyl-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-2-naphthylacetamide,  
2-(7-fluoro-1H-benzimidazol-1-yl)-N-2-naphthylacetamide,  
2-(7-cyano-1H-benzimidazol-1-yl)-N-(4-methoxy-2-naphthyl)acetamide,  
15 2-(7-fluoro-1H-benzimidazol-1-yl)-N-(4-methoxy-2-naphthyl)acetamide,  
N-[3-methoxy-5-(tetrahydro-2H-pyran-2-ylmethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-(2-isopropoxyethoxy)-5-methoxyphenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
20 N-[3,5-bis(2-ethoxyethoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-methoxy-5-[2-(2-oxopyrrolidin-1-yl)ethoxy]phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-methoxy-5-(3-morpholin-4-ylpropoxy)phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
25 N,N-diethyl-2-(3-methoxy-5-[(7-nitro-1H-benzimidazol-1-yl)acetyl]amino)phenoxyacetamide,  
N-[3-methoxy-5-[(1-methylpiperidin-2-yl)methoxy]phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,  
N-[3-[2-(1H-imidazol-1-yl)ethoxy]-5-methoxyphenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide, and  
30 N-[3-methoxy-5-[(1-methyl-1H-imidazol-5-yl)methoxy]phenyl]-2-(7-nitro-1H-benzimidazol-1-yl)acetamide,

or salts, solvates or solvated salts thereof.

5 5. A pharmaceutical composition comprising as active ingredient a therapeutically effective amount of the compound according to claims 1, 2, 3 or 4, in association with one or more pharmaceutically acceptable diluents, excipients and/or inert carriers.

6. The pharmaceutical composition according to claim 5, for use in the treatment of VR1 mediated disorders and for treatment of acute and chronic pain disorders, acute and chronic neuropathic pain and acute and chronic inflammatory pain, and respiratory diseases.

10 7. The compound according to claims 1, 2, 3 or 4, for use as a medicament.

8. Use of a compounds according to claims 1, 2, 3 or 4, in the manufacture of a medicament for treatment of VR1 mediated disorders.

15 9. The use according to claim 8 for treatment of acute and chronic pain disorders.

10. The use according to claim 8 for treatment of acute and chronic neuropathic pain.

20 11. The use according to claim 8 for treatment of acute and chronic inflammatory pain.

12. The use according to claim 8 for treatment of low back pain, post-operative pain, visceral pains like chronic pelvic pain, cystitis, including interstitial cystitis and pain related thereto, ischeamic, sciatica, diabetic neuropathy, multiple sclerosis, arthritis, fibromyalgia, psoriasis, cancer, emesis, urinary incontinence, hyperactive bladder, HIV neuropathy, gastro-esophageal reflux disease (GERD), irritable bowel syndrome (IBS), inflammatory  
25 bowel disease (IBD) and pancreatitis.

13. The use according to claim 8 for treatment of respiratory diseases.

30 14. A method of treatment of VR1 mediated disorders and for treatment of acute and chronic pain disorders, acute and chronic neuropathic pain and acute and chronic inflam-

matory pain, and respiratory diseases, comprising administering to a mammal, including man in need of such treatment, a therapeutically effective amount of a compound according to claims 1, 2, 3 or 4.

- 5 15. The compound selected from the group consisting of  
3-methoxy-5-(methoxymethyl)aniline,  
3-(methoxymethyl)-5-(trifluoromethyl)aniline,  
1-(methoxymethyl)-3-nitro-5-(trifluoromethyl)benzene,  
1-[3-amino-5-(trifluoromethyl)phenyl]ethanone,  
10 (7-chloro-6-methoxy-1H-benzimidazol-1-yl)acetic acid,  
2-[(2-chloro-3-methoxy-6-nitrophenyl)amino]ethanol,  
2-(7-chloro-6-methoxy-1H-benzimidazol-1-yl)ethanol,  
3-(2-methoxyethoxy)-5-(trifluoromethyl)aniline,  
1-(2-methoxyethoxy)-3-nitro-5-(trifluoromethyl)benzene,  
15 3-methoxy-5-(tetrahydrofuran-2-ylmethoxy)aniline,  
2-[(3-methoxy-5-nitrophenoxy)methyl]tetrahydrofuran,  
3-methoxy-5-(tetrahydrofuran-3-yloxy)aniline,  
3-(3-methoxy-5-nitrophenoxy)tetrahydrofuran,  
5,6-dihydro-4H-imidazo[4,5,1-ij]quinoline-4-carboxylic acid,  
20 methyl 8-amino-1,2,3,4-tetrahydroquinoline-2-carboxylate,  
(7-pyridin-2-yl-1H-benzimidazol-1-yl)acetic acid,  
methyl (7-bromo-1H-benzimidazol-1-yl)acetate,  
methyl (7-pyridin-2-yl-1H-benzimidazol-1-yl)acetate  
3-methoxy-5-(tetrahydro-2H-pyran-2-ylmethoxy)aniline, and  
25 3-(2-isopropoxyethoxy)-5-methoxyaniline


16. Use of the compounds according to claim 15 as intermediates in the preparation of compounds according to claims 1, 2, 3 or 4.

## PATENT COOPERATION TREATY

## PCT

## INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)



Applicant's or agent's file reference <b>101543-1 WO</b>	FOR FURTHER ACTION <small>see Form PCT/ISA/220 as well as, where applicable, item 5 below.</small>	
International application No. <b>PCT/SE 2005/001364</b>	International filing date (day/month/year) <b>19 Sept. 2005</b>	(Earliest) Priority Date (day/month/year) <b>21 Sept. 2004</b>
Applicant <b>AstraZeneca AB et al</b>		

This international search report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This international search report consists of a total of 6 sheets.

☐ It is also accompanied by a copy of each prior art document cited in this report.

## 1. Basis of the report

a. With regard to the language, the international search was carried out on the basis of:

- ☒ the international application in the language in which it was filed  
☐ a translation of the international application into \_\_\_\_\_, which is the language of a translation furnished for the purposes of international search (Rules 12.3(a) and 23.1(b))

b. ☐ With regard to any nucleotide and/or amino acid sequence disclosed in the international application, see Box No. I.

2. ☒ Certain claims were found unsearchable (see Box No. II)

3. ☒ Unity of invention is lacking (see Box No. III)

4. With regard to the title,

- ☒ the text is approved as submitted by the applicant.  
☐ the text has been established by this Authority to read as follows:

5. With regard to the abstract,

- ☒ the text is approved as submitted by the applicant.  
☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box No. IV. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. With regard to the drawings,

- a. the figure of the drawings to be published with the abstract is Figure No. —  
☐ as suggested by the applicant.  
☐ as selected by this Authority, because the applicant failed to suggest a figure.  
☐ as selected by this Authority, because this figure better characterizes the invention.
- b. ☐ none of the figures is to be published with the abstract.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/SE 2005/001364

## A. CLASSIFICATION OF SUBJECT MATTER

IPC: see extra sheet

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-INTERNAL, WPI DATA, PAJ, CA

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	STN International, File REGISTRY, see RN 743444-08-08, 13 Sep 2004 --	1-14
P,X	WO 2004100865 A2 (ASTRAZENECA AB), 25 November 2004 (25.11.2004), see examples --	1-14
X	US 20040092569 A1 (DEMAINE ET AL), 13 May 2004 (13.05.2004), treatment of pain and inflammations, examples 1-8, abstract --	1,5-14
A	US 20030149050 A1 (JAGTAP ET AL), 7 August 2003 (07.08.2003), inhibitors of inflammation, column 1, example 19 (bensimidazole) --	1-8,11,14

☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.

\* Special categories of cited documents

"A" document defining the general state of the art which is not considered to be of particular relevance  
"E" earlier application or patent but published on or after the international filing date  
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reasons (as specified)  
"O" document referring to an oral disclosure, use, exhibition or other means  
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  
"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone  
"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art  
"&" document member of the same patent family

Date of the actual completion of the international search

2 February 2006

Date of mailing of the international search report

03-02-2006

Name and mailing address of the ISA/  
Swedish Patent Office  
Box 5055, S-102 42 STOCKHOLM  
Facsimile No. +46 8 666 02 86

Authorized officer

Eva Johansson/EÖ  
Telephone No. +46 8 782 25 00

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 2005/001364

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	B. VIJAYA KUMAR et al, "Cyanoethylation of Benzimidazoles: Synthesis & Biological Activities of Some New 1-(beta-Cyanoethyl)benzimidazoles & Their Derivatives", Indian Journal of Chemistry, October 1985, Vol. 24B, p.1098-1101 treatment of pain and inflammations  --	1-14
A	WO 0196336 A2 (WARNER-LABERT COMPANY), 20 December 2001 (20.12.2001), treatment of inflammation, abstract, schemes 12 and 13  --	1-8,11,14
A	EP 0419210 A1 (PFIZER INC.), 27 March 1991 (27.03.1991), benzimidazoles as antiinflammatory agents  --	1-8,11,14
A	US 3624103 A (FRANDO E MARTIIS), 30 November 1971 (30.11.1971), anti-inflammatory and analgesic effects, abstract  --	1-8,11,14
A	US 20030158188 A1 (LEE ET AL), 21 August 2003 (21.08.2003), VRI antagonists  --	1-14
A	US 20040152690 A1 (BLAN ET AL), 5 August 2004 (05.08.2004), treatment of inflammation and pain, abstract  -- -----	1-14



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/SE2005/001364**Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 14  
because they relate to subject matter not required to be searched by this Authority, namely:  
Claim 14 relate to a method of treatment of the human body by therapy /Rule 39.1(iv). Nevertheless, a search has been executed for this claim. The search has been based on the alleged effects of the compounds.
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

In order to fulfil the requirements of unity of invention, it is necessary that the intermediate compounds are closely interconnected with the end products. Such close connection requires that the essential structural part of the end product is incorporated by the intermediate compound. However, the

.../...

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.: 1 - 14

**Remark on Protest**

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.  
PCT/SE2005/001364

**Box III**

present application lacks a single general inventive concept based on the above principle. This leads to the presence of the subjects listed below, each falling under its own restricted inventive concept.

I: Claim 1-14 concerning final compounds.

II: Claims 15 and 16 concerning the intermediate compounds that are anilines or nitrophenyls, i.e. compounds 1-4, 6, 8-13, 15 and 19-20 of claim 15.

III: 15 and 16 concerning the intermediate compounds comprising benzimidazole, i.e. compounds 5, 7, 14 and 16-18 of claim 15.

A partial search has been carried out, which relates to the invention I mentioned above.

The present application has been considered to contain 3 inventions which are not linked such that they form a single general inventive concept, as required by Rule 13 PCT.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/SE2005/001364

**INTERNATIONAL PATENT CLASSIFICATION (IPC):**

**C07D 401/04** (2006.01)  
**A61K 31/4184** (2006.01)  
**A61K 31/4439** (2006.01)  
**A61P 29/00** (2006.01)  
**C07D 235/06** (2006.01)  
**C07D 405/12** (2006.01)

INTERNATIONAL SEARCH REPORT  
Information on patent family members

26/11/2005

International application No.  
PCT/SE 2005/001364

WO	2004100865	A2	25/11/2004	AU	2003261053 A	00/00/0000
				CA	2498151 A	01/04/2004
				EP	1543094 A	22/06/2005
				SE	0301446 D	00/00/0000
				SE	0400043 D	00/00/0000
US	20040092569	A1	13/05/2004	NONE		
US	20030149050	A1	07/08/2003	US	6903079 B	07/06/2005
				AU	4991801 A	23/10/2001
				US	6534651 B	18/03/2003
				US	20020095044 A	18/07/2002
				WO	0177075 A	18/10/2001
WO	0196336	A2	20/12/2001	AU	5970701 A	24/12/2001
				BR	0111544 A	01/07/2003
				CA	2412462 A	20/12/2001
				EP	1294718 A	26/03/2003
				JP	2004503553 T	05/02/2004
				MX	PA02010763 A	10/03/2003
				US	6943174 B	13/09/2005
				US	20040014759 A	22/01/2004
EP	0419210	A1	27/03/1991	SE	0419210 T3	
				AT	113586 T	15/11/1994
				CA	2025849 A	23/03/1991
				DE	69013836 D, T	23/03/1995
				DK	419210 T	09/01/1995
				ES	2062395 T	16/12/1994
				FI	904659 D	00/00/0000
				IE	63816 B	14/06/1995
				IE	903408 A	10/04/1991
				JP	1876018 C	07/10/1994
				JP	3109378 A	09/05/1991
				JP	6000759 B	05/01/1994
				PT	95371 A	22/05/1991
				US	5141950 A	25/08/1992
US	3624103	A	30/11/1971	BE	810092 A	16/05/1974
				IT	1036004 B	30/10/1979
				NL	6810284 A	25/11/1969
US	20030158188	A1	21/08/2003	CA	2476936 A	28/08/2003
				EP	1478363 A	24/11/2004
				WO	03070247 A	28/08/2003
US	20040152690	A1	05/08/2004	AU	2003301436 A	00/00/0000
				CA	2501539 A	29/04/2004
				EP	1551811 A	13/07/2005
				WO	2004035549 A	29/04/2004